

# Marine hypoxia/anoxia as a source of CH<sub>4</sub> and N<sub>2</sub>O

S. W. A. Naqvi<sup>1,2</sup>, H. W. Bange<sup>3</sup>, L. Farías<sup>4</sup>, P. M. S. Monteiro<sup>5</sup>, M. I. Scranton<sup>6</sup>, and J. Zhang<sup>7</sup>

<sup>1</sup>National Institute of Oceanography (Council of Scientific & Industrial Research), Dona Paula, Goa 403 004, India

<sup>2</sup>Max-Planck Institut für Marine Mikrobiologie, Celsiusstrasse 1, 28359 Bremen, Germany

<sup>3</sup>Forschungsbereich Marine Biogeochemie, IFM-GEOMAR, Düsternbrooker Weg 20, 24105 Kiel, Germany

<sup>4</sup>Laboratorio de Procesos Oceanográficos y Clima (PROFC), Departamento de Oceanografía y Centro de Investigación Oceanográfica en el Pacífico Suroriental (COPAS), Universidad de Concepción, Casilla 160-C, Concepción, Chile

<sup>5</sup>Department of Oceanography, University of Cape Town, Rondebosch, South Africa

<sup>6</sup>School of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook NY 11794, USA

<sup>7</sup>State Key Laboratory of Estuarine and Coastal Research, East China Normal University, 3663 Zhongshan Road North, 200062 Shanghai, China

Received: 28 August 2009 – Published in Biogeosciences Discuss.: 2 October 2009

Revised: 21 June 2010 – Accepted: 22 June 2010 – Published: 12 July 2010

**Abstract.** We review here the available information on methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) from major marine, mostly coastal, oxygen (O<sub>2</sub>)-deficient zones formed both naturally and as a result of human activities (mainly eutrophication). Concentrations of both gases in subsurface waters are affected by ambient O<sub>2</sub> levels to varying degrees. Organic matter supply to seafloor appears to be the primary factor controlling CH<sub>4</sub> production in sediments and its supply to (and concentration in) overlying waters, with bottom-water O<sub>2</sub>-deficiency exerting only a modulating effect. High (micromolar level) CH<sub>4</sub> accumulation occurs in anoxic (sulphidic) waters of silled basins, such as the Black Sea and Cariaco Basin, and over the highly productive Namibian shelf. In other regions experiencing various degrees of O<sub>2</sub>-deficiency (hypoxia to anoxia), CH<sub>4</sub> concentrations vary from a few to hundreds of nanomolar levels. Since coastal O<sub>2</sub>-deficient zones are generally very productive and are sometimes located close to river mouths and submarine hydrocarbon seeps, it is difficult to differentiate any O<sub>2</sub>-deficiency-induced enhancement from in situ production of CH<sub>4</sub> in the water column and its inputs through freshwater runoff or seepage from sediments. While the role of bottom-water O<sub>2</sub>-deficiency in CH<sub>4</sub> formation appears to be secondary, even when CH<sub>4</sub> accumulates in O<sub>2</sub>-deficient subsurface waters, methanotrophic activity severely restricts its diffusive efflux to the atmosphere. As a result, an intensification or expansion of coastal O<sub>2</sub>-deficient zones will probably

not drastically change the present status where emission from the ocean as a whole forms an insignificant term in the atmospheric CH<sub>4</sub> budget. The situation is different for N<sub>2</sub>O, the production of which is greatly enhanced in low-O<sub>2</sub> waters, and although it is lost through denitrification in most suboxic and anoxic environments, the peripheries of such environments offer most suitable conditions for its production, with the exception of enclosed anoxic basins. Most O<sub>2</sub>-deficient systems serve as strong net sources of N<sub>2</sub>O to the atmosphere. This is especially true for coastal upwelling regions with shallow O<sub>2</sub>-deficient zones where a dramatic increase in N<sub>2</sub>O production often occurs in rapidly denitrifying waters. Nitrous oxide emissions from these zones are globally significant, and so their ongoing intensification and expansion is likely to lead to a significant increase in N<sub>2</sub>O emission from the ocean. However, a meaningful quantitative prediction of this increase is not possible at present because of continuing uncertainties concerning the formative pathways to N<sub>2</sub>O as well as insufficient data from key coastal regions.

## 1 Introduction

Methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) are two important trace constituents of the atmosphere. Both CH<sub>4</sub> and N<sub>2</sub>O are potent greenhouse gases that are approximately 300 and 25 times more effective, respectively, on a per molecule basis than carbon dioxide (CO<sub>2</sub>) in trapping heat. They also play important roles in atmospheric chemistry – CH<sub>4</sub> is the most abundant hydrocarbon in the atmosphere and N<sub>2</sub>O is an important precursor for nitric oxide (NO) radicals which



Correspondence to: S. W. A. Naqvi  
(naqvi@nio.org)

**Table 1.** Definitions of various stages of O<sub>2</sub> deficiency.

Stage	Criteria	Remarks
Hypoxia	$0.1 < \text{O}_2$ (Winkler) $\leq 1.4 \text{ ml l}^{-1}$ (see footnote <sup>a</sup> ); $\text{NO}_2^- = 0 \mu\text{M}$ , $\text{NO}_3^- > 0 \mu\text{M}$	The upper limit is based on physiological consideration. Oxygen concentrations below $\sim 2 \text{ mg l}^{-1}$ induce avoidance, or altered behavior, growth, reproduction or survivorship of many marine organisms (Levin et al., 2009, and references therein)
Suboxia	$\text{O}_2 \leq 0.1 \text{ ml l}^{-1}$ , $\text{NO}_2^-$ , $\text{NO}_3^- > 0 \mu\text{M}$ ; $\text{H}_2\text{S} = 0 \mu\text{M}$	The actual O <sub>2</sub> levels are probably in the nM range (Revsbech et al., 2009). This allows reduction of elements such as N, I, Mn and Fe (but not S) with denitrification being the dominant respiratory process.
Anoxia	$\text{O}_2 = 0 \text{ ml l}^{-1}$ , $\text{NO}_2^-$ , $\text{NO}_3^- = 0 \mu\text{M}$ , $\text{H}_2\text{S} > 0 \mu\text{M}$	The environment is completely anaerobic with sulphate reduction being the dominant respiratory process.

<sup>a</sup>  $1 \text{ ml l}^{-1} = 1.43 \text{ mg l}^{-1} = 44.64 \mu\text{M}$

are involved in the destruction of ozone in the stratosphere. Both gases are produced by natural processes as well as by human activities. The latter include landfills, livestock, fossil fuel exploitation, agriculture (rice paddies) and wastewater treatment for CH<sub>4</sub>, and agriculture (application of N-fertilizers) and fossil fuel combustion for N<sub>2</sub>O. As a result of anthropogenic emissions, average atmospheric concentrations of CH<sub>4</sub> and N<sub>2</sub>O by 2007 had risen by 156% and 19% above the pre-industrial levels to 1789 and 321 parts per billion (ppb), respectively (Anonymous, 2008).

The ocean serves as a large source of N<sub>2</sub>O to the atmosphere, accounting for at least one third of all natural emissions (IPCC, 2007; Bange, 2006a). Although the ocean also emits CH<sub>4</sub> to the atmosphere, its contribution to the atmospheric CH<sub>4</sub> budget is minor (<2% – Reeburgh, 2007). Unlike the terrestrial sources, the impingement of human activities on oceanic emissions of these gases is not well understood and poorly quantified. One important factor that exerts the key control on biological cycling of these gases in the ocean is the redox state of the environment, which is determined by the ambient oxygen (O<sub>2</sub>) concentration. Containing carbon in its most reduced (−4) form, CH<sub>4</sub> is produced in significant quantities only in anoxic environments. In the case of N<sub>2</sub>O (which contains nitrogen in an intermediate oxidation state of +1), both oxidative (nitrification) and reductive (denitrification) production pathways exist. Denitrification [reduction of nitrate (NO<sub>3</sub><sup>−</sup>) to elemental nitrogen (N<sub>2</sub>) with N<sub>2</sub>O as an intermediate] is, of course, an anaerobic process. But even in nitrification [oxidation of ammonium (NH<sub>4</sub><sup>+</sup>) to NO<sub>3</sub><sup>−</sup> where N<sub>2</sub>O is formed as a byproduct], the yield of N<sub>2</sub>O relative to NO<sub>3</sub><sup>−</sup> increases as the O<sub>2</sub> concentrations fall below about  $0.5 \text{ ml l}^{-1}$  ( $\sim 22 \mu\text{M}$ ) (Goreau et al., 1980). Thus, changes in O<sub>2</sub> distribution may alter source strengths of CH<sub>4</sub> and N<sub>2</sub>O. Such changes in subsurface O<sub>2</sub> field may be forced by altered circulation/stratification in re-

sponse to greenhouse warming, and/or by elevated respiration of organic matter produced as a result of enhanced nutrient supply from land.


The extent to which human activities are affecting/have affected physical processes that control subsurface O<sub>2</sub> distribution is unclear. It has been suggested that coastal upwelling has become more vigorous since the 1940s due to an intensification of continental thermal lows adjacent to the eastern boundary upwelling regions off California, Northwest Africa, Iberian Peninsula and Peru (Bakun, 1990). More recently, the observed intensification of hypoxic conditions (see Table 1 for the terms used in this paper to define various stages of O<sub>2</sub> deficiency) in the California Current region has been attributed to a change in water circulation over the shelf (Grantham et al., 2004; Bograd et al., 2008; Chan et al., 2008). Moreover, the observed and modelled decreases in the O<sub>2</sub> concentration in subsurface waters including an expansion of oceanic oxygen minimum zones (OMZs) point to physical causes such as decrease in surface concentration and slower subsurface ventilation (Joos et al., 2003; Stramma et al., 2008; Oschlies et al., 2008; Shaffer et al., 2009). The growing deposition of anthropogenic nitrogen from the atmosphere that extends well beyond coastal waters (Galloway et al., 2004; Jickels, 2006; Duce et al., 2008) is also expected to contribute to the ongoing subsurface O<sub>2</sub> decline in the open ocean. According to Duce et al. (2008), atmospheric deposition of anthropogenic nitrogen could have led to an increase in oceanic N<sub>2</sub>O emission by  $\sim 1.6 \text{ Tg N a}^{-1}$  ( $T = 10^{12}$ ). While the exact mechanism of this increase is not known, some part of it might be due to enhanced production associated with O<sub>2</sub> depletion. In contrast with the atmospheric deposition, the riverine supply of reactive nitrogen that has been greatly altered by human activities (Seitzinger et al., 2002; Galloway et al., 2004) is largely confined to the coastal zone. Moreover, while the atmospheric deposition is mostly

restricted to nitrogen, rivers also bring other macronutrients (phosphorus and silicon), of which phosphorus flux has also more than doubled as a result of human activities (Smith et al., 2003). The enhanced supply of (anthropogenic) nutrients (eutrophication) to coastal regions has led to development of hypoxia in many parts of the world (Diaz and Rosenberg, 2008, and references therein) that would, among other things, affect the production of CH<sub>4</sub> and N<sub>2</sub>O. Here we review the information available on these gases from marine systems that are affected by natural as well as human-induced O<sub>2</sub> deficiency. We focus on major coastal O<sub>2</sub>-deficient zones, as well as semi-enclosed basins, while a large number of smaller water bodies (estuaries, lagoons and fjords) that also experience hypoxia on a variety of time scales (Diaz and Rosenberg, 2008; Kemp et al., 2009; Rabalais et al., 2009) are not considered.

## 2 Processes responsible for formation of O<sub>2</sub>-depleted coastal systems

Oxygen depletion in seawater, as in all other aquatic systems, can occur due to natural processes as well as anthropogenic factors, and it is sometimes difficult to de-convolve the effects of the two forcings. For example, controversy persists concerning the cause(s) of hypoxia even for the best known of all coastal hypoxic systems – the “dead zone” of the Louisiana Shelf in the northern Gulf of Mexico (Rabalais et al., 2007; Swarzenski et al., 2008). All natural O<sub>2</sub>-deficient aquatic environments have arguably been affected by human activities to varying degrees. Nonetheless, it is possible in most cases to identify the dominant driver of hypoxia. Thus, out of the systems being examined here (Fig. 1), hypoxia in the East China Sea, Chesapeake Bay, Gulf of Mexico and Tokyo Bay is largely human-induced, whereas in the remaining regions it is primarily of natural origin.

The most important oceanic O<sub>2</sub>-deficient environments are formed naturally and have existed with varying intensities through geological times (Neretin, 2006). Today, a number of semi-enclosed/land-locked seas and fjords experience anoxia due to stagnation of subsurface waters. The Black Sea, the Cariaco Basin and the central Baltic Sea are the best known of such water bodies. Of these, the first two are permanently anoxic below their sill depths whereas the third is periodically so. By contrast, in the open ocean, severe O<sub>2</sub> depletion only occurs within the mesopelagic realm in a few well-demarcated geographical areas (Deuser, 1975). Such oxygen minimum zones (OMZs) can be easily identified in global maps of O<sub>2</sub> at depths of few hundred meters, an example of which (for 150 m, close to the upper boundary of OMZs) is shown in Fig. 1. The OMZs are located in the tropics and subtropics along the eastern boundaries of the Atlantic and Pacific oceans. They develop because of a combination of slower thermocline water renewal than in the subtropical gyres (which is why they are sometimes referred



**Fig. 1.** Mean annual dissolved O<sub>2</sub> concentration ( $\mu\text{mol kg}^{-1}$ ) at 150 m depth.  $1^\circ \times 1^\circ$  gridded O<sub>2</sub> data were taken from World Ocean Atlas 2005 (Garcia et al., 2006) and converted to  $\mu\text{mol kg}^{-1}$ . The O<sub>2</sub>-depleted zones discussed in the text are marked: 1, Off SW Africa; 2, Baltic Sea; 3, Black Sea; 4, Arabian Sea; 5, Bay of Bengal; 6, East China Sea; 7, Tokyo Bay; 8, Saanich Inlet; 9, Eastern North Pacific; 10, Gulf of Mexico; 11, Chesapeake Bay; 12, Eastern South Pacific; 13, Cariaco Basin; 14, Off NW Africa. Squares depict semi-enclosed basins that experience sulphate reduction whereas circles/ovals depict other low O<sub>2</sub>/hypoxic/suboxic (including seasonally anoxic) environments along open coasts.

to as “shadow zones”) (Luyten et al., 1983) and higher O<sub>2</sub> demand for respiration arising from high biological production in the overlying surface waters fuelled by nutrient enrichment through upwelling (Gruber, 2004). The OMZs in the Pacific (off western North America – Mexico, and off western South America – Chile and Peru) are more voluminous than those in the Atlantic (off Southwest Africa – Namibia, and off Northwest Africa – Mauritania) because of the generally lower subsurface O<sub>2</sub> concentrations throughout the Pacific. In all but one (off Northwest Africa) of these four eastern-boundary upwelling ecosystems (EBUEs), O<sub>2</sub> concentrations fall to suboxic levels. Due to its unusual geography – its northward expanse is limited by the South Asian land mass at low latitudes – and resultant unique monsoonal circulation, the eastern boundary of the Indian Ocean does not experience vigorous upwelling and associated O<sub>2</sub> deficiency. Instead, the dominant variability in O<sub>2</sub> below the surface layer occurs in the north-south direction (Wyrki, 1971). The OMZ is particularly intense ( $\text{O}_2 < 0.02 \text{ ml l}^{-1}$ ,  $\sim 1 \mu\text{M}$ ) and thick (between  $\sim 100/150$  and  $1200 \text{ m}$  – Codispoti et al., 2001) in the Northwest Indian Ocean (Arabian Sea), which is the most productive part of the Indian Ocean because of nutrient enrichment of the euphotic zone through convective mixing in winter and upwelling in summer (Naqvi et al., 2003). Extension of offshore OMZs to the coastal region occurs through the process of upwelling. Thus, advection of low-O<sub>2</sub> waters is prerequisite to (natural) development of coastal hypoxia. However, in addition to the initial O<sub>2</sub>

content of upwelling water, intensity of O<sub>2</sub> deficiency over the shelf is modulated by a number of factors – local circulation/hydrography especially stratification, upwelling intensity, shelf width, and primary production – that together determine the balance between oxygen consumption and supply in subsurface waters.

Coastal hypoxic zones produced due to anthropogenic activities, whose number is steadily increasing over the past few decades (Diaz and Rosenberg, 2008), are mostly located in nearshore waters and estuaries in areas that are subject to high loading of nutrients and/or organic matter from terrestrial sources. Most of this loading occurs through river runoff, although in the case of nitrogen atmospheric inputs are also important (Duce et al., 2008). This in conjunction with stratification caused by freshwater additions makes estuaries and coastal zones off major rivers especially susceptible to deoxygenation. The continental shelf off the Mississippi River mouth in the Gulf of Mexico, the Chesapeake Bay that receives outflows from a number of rivers such as the Susquehanna and the Potomac, and the coastal region off the Changjiang in the East China Sea (ECS) are the best examples of such systems. Hypoxia in these systems generally exhibits large seasonality due to changes in river runoff and solar insolation, and is generally at its maximum intensity in summer (e.g. Rabalais and Turner, 2006).

### 3 Brief overview of CH<sub>4</sub> and N<sub>2</sub>O cycling in the ocean

#### 3.1 Methane

The oceanic CH<sub>4</sub> biogeochemistry has recently been reviewed by Reeburgh (2007). Its concentrations in the open-ocean water column are generally quite low (a few nanomolar). Large CH<sub>4</sub> build-up does not occur even in open-ocean suboxic zones because methanogenesis, microbial production of CH<sub>4</sub> from CO<sub>2</sub> or acetate, is inhibited by the presence of other electron accepters such as oxygen and sulphate. Conditions favouring this process usually develop in sediments where below a certain horizon (the sulphate-CH<sub>4</sub> transition) sulphate is almost totally depleted. Despite very high (millimolar) CH<sub>4</sub> concentrations below the sulphate-CH<sub>4</sub> interface, very low levels in the water column – the deep water is typically undersaturated with respect to atmospheric CH<sub>4</sub> – point to efficient oxidation, both in the sediment and water column (Rehder et al., 1999; Reeburgh, 2007).

The upper ocean is characterized by relative higher CH<sub>4</sub> concentrations – close to saturation at the sea-surface with a maximum located near the base of the mixed layer. The cause of the ubiquitous occurrence of the maximum in well-oxygenated water – the so-called oceanic CH<sub>4</sub> paradox – has long been debated. Several explanations for the subsurface CH<sub>4</sub> maximum have been proposed and include both in situ production and advective transport of CH<sub>4</sub> produced elsewhere. For example, it has been speculated that production

within anoxic environments, most probably within guts of zooplankton, accounts for the maximum (Oremland, 1979; Reeburgh, 2007, and references therein). Coastal waters may also contain elevated CH<sub>4</sub> concentrations due to supply from marshlands/estuaries (Scranton and McShane, 1991; Bange et al., 1994; Jayakumar et al., 2001; Bange, 2006b) and from sediments, particularly in areas of hydrocarbon seepage or hydrate destabilization, resulting in pronounced subsurface maxima extending offshore from the continental margin that are distinct from the mixed layer maximum mentioned above (e.g., Scranton and Brewer, 1977; Brooks et al., 1981; Cynar and Yayanos, 1991; Ward, 1992; Tilbrook and Karl, 1995). More recently, it has been shown that CH<sub>4</sub> may be produced aerobically through decomposition of methylphosphonate that may serve as a source of phosphorus in phosphate-poor environments such as tropical surface waters (Karl et al., 2008). Subsequently, Damm et al. (2010) reported CH<sub>4</sub> production in surface waters of the central Arctic Ocean that contained sufficient phosphate but were nitrate-depleted. These authors proposed that under phosphate-replete conditions methylated compounds like dimethylsulfoniopropionate (DMSP) could be utilized by bacteria as the source of carbon. Although such a production of CH<sub>4</sub> as a byproduct of bacterial metabolism in aerobic waters has been shown to be thermodynamically feasible, the paradox still needs to be fully resolved.

#### 3.2 Nitrous oxide

Bange (2008) has provided a comprehensive review of oceanic biogeochemistry of N<sub>2</sub>O. Surface waters of the ocean are generally slightly supersaturated with respect to atmospheric N<sub>2</sub>O (~4%; Nevison et al., 1995), but much higher supersaturation occurs in regions that experience coastal upwelling and are affected by subsurface O<sub>2</sub>-deficiency (e.g. Bange et al., 2001a; Nevison et al., 2004). An important aspect of N<sub>2</sub>O distribution in the ocean is the inverse relationship with O<sub>2</sub>; this feature, recognized by early workers, prompted the suggestion that nitrification is the dominant process responsible for N<sub>2</sub>O production (Yoshinari, 1976). Subsequently several authors reported linear relationships between  $\Delta\text{N}_2\text{O}$  ( $=\text{N}_2\text{O}_{\text{observed}}-\text{N}_2\text{O}_{\text{saturation}}$ ) and apparent oxygen utilization ( $\text{AOU}=\text{O}_2_{\text{saturation}}-\text{O}_2_{\text{observed}}$ ) from various oceanic areas (e.g. Elkins et al., 1978; Oudot et al., 1990; Naqvi and Noronha, 1991; Naqvi et al., 1994). This relationship holds as long as the O<sub>2</sub> concentration does not fall below ~0.5 ml l<sup>-1</sup> after which a sharp non-linear N<sub>2</sub>O build-up occurs as O<sub>2</sub> is further consumed (Codispoti and Christensen, 1985; Codispoti et al., 1992). Once the system turns suboxic, however, a depletion of N<sub>2</sub>O takes place to levels below saturation, a trend consistently seen in all major open-ocean suboxic zones (Elkins et al., 1978; Cohen and Gordon, 1978; Naqvi and Noronha, 1991; Farias et al., 2009a). The sensitivity of N<sub>2</sub>O production and consumption to minor changes in O<sub>2</sub> in the low range has led to a

large number of studies focussing on N<sub>2</sub>O distribution in the OMZs (Elkins et al., 1978; Cohen and Gordon, 1978; Codispoti and Christensen, 1985; Codispoti et al., 1992; Law and Owens, 1990; Naqvi and Noronha, 1991; Upstill-Goddard et al., 1999; Bange et al., 2001b; Farias and Cornejo, 2007; Cornejo et al., 2006, 2007; Farias et al., 2009a). The low N<sub>2</sub>O concentrations within nitrite (NO<sub>2</sub><sup>-</sup>)-bearing waters are widely believed to arise from its utilization as an electron acceptor by heterotrophic bacteria (which convert it to N<sub>2</sub>). Despite the existence of a sink for N<sub>2</sub>O within the secondary nitrite maximum of open-ocean OMZs, these zones still serve as disproportionately large net sources of N<sub>2</sub>O because of high N<sub>2</sub>O concentration at their peripheries. The mechanisms responsible for this accumulation still continue to be poorly understood. Based on the isotopic evidence, Dore et al. (1998) suggested that the shallow N<sub>2</sub>O maximum in the North Pacific is formed largely through nitrification, whereas Naqvi et al. (1998) proposed that a coupling between nitrification and denitrification could be a more important formative pathway for the upper N<sub>2</sub>O maximum in the Arabian Sea. Nicholls et al. (2006) demonstrated that this N<sub>2</sub>O maximum is produced largely through reduction of NO<sub>2</sub><sup>-</sup> to N<sub>2</sub>O, but whether this reduction is carried out by denitrifiers or nitrifiers still needs to be resolved (see also Shailaja et al., 2006). Anaerobic ammonium oxidation (anammox) and dissimilatory nitrate reduction to ammonium (DNRA) are two other potential processes that can produce N<sub>2</sub>O from NO<sub>2</sub><sup>-</sup>. However, in laboratory cultures anammox bacteria have not been found to produce significant amounts of N<sub>2</sub>O (van der Star, 2008) and, except for one study (Lam et al., 2009), DNRA has not been considered to be a major process in the oceanic water column. In an intertidal seagrass meadow, though, DNRA by fermentative and/or sulphate-reducing bacteria was proposed to serve as a source of N<sub>2</sub>O (Welsh et al., 2001). As we will see in the following section, the pattern of variability observed in coastal suboxic zones sometimes deviates from the trend observed in open-ocean OMZs.

## 4 Methane and nitrous oxide in major marine O<sub>2</sub>-deficient zones other than enclosed anoxic basins

### 4.1 Eastern Atlantic Ocean

#### 4.1.1 Off Northwest Africa

The region off Northwest Africa (Morocco-Mauritania) in the eastern equatorial Atlantic Ocean (the Canary Current System) is one of the four major EBUEs. It is also the only one where subsurface O<sub>2</sub> depletion does not qualify to be classified as hypoxic, because of the relatively high initial O<sub>2</sub> content of waters upwelling over the shelf. However, given the ongoing deoxygenation of subsurface waters in the North Atlantic, there are concerns that hypoxia in subsurface wa-

ters may develop in future in this region as well (Stramma et al., 2008).

Kock et al. (2008) carried out extensive measurements of CH<sub>4</sub> in atmospheric and surface seawater samples collected off Mauritania during two cruises conducted in March/April 2005 and February 2007. They found saturations reaching up to 200% in the upwelled water. The annual CH<sub>4</sub> emission from the region was quantified as 1.6–2.9 Gg CH<sub>4</sub>, which is not very significant in comparison to global supply to the atmosphere.


There are at present no published data on N<sub>2</sub>O from this area. However, in October–November 2002, Walter et al. (2004) measured N<sub>2</sub>O in surface waters along a transect at 10° N. At the eastern end of the transect, over the African shelf to the south of the Mauritanian upwelling zone, N<sub>2</sub>O saturation reached a maximum of 113%, which was ascribed to N<sub>2</sub>O supply by river or ground waters.

#### 4.1.2 Off Southwest Africa

Upwelling off Southwest Africa (Namibia) supports the highest primary production rates of all the four major EBUEs (Carr, 2002), and degradation of copious organic matter in the water column and shelf sediments culminates in the occurrence of sulphidic conditions in bottom waters extending offshore to the shelf edge off the Walvis Bay (Chapman and Shannon, 1985; Brüchert et al., 2006, 2009; van der Plas et al., 2007; Lavik et al., 2009). In addition to respiration of locally-produced organic matter, the intensity of O<sub>2</sub> deficiency is also modulated by the initial O<sub>2</sub> content of upwelling waters. There are two main upwelling centres in this region located off Cape Frio in the north and Luderitz in the south. Upwelling at these locations is fed by intermediate waters with very different O<sub>2</sub> contents: the hypoxic Angola Basin Central Water off Cape Frio and more oxygenated Cape Basin South Atlantic Central Water off Luderitz, with the boundary between the two subsurface water types located around 25° S latitude (Monteiro et al., 2008). Once over the shelf, these waters move northward (from Luderitz) or southward (from Cape Frio) and it is the relative contribution from these sources that controls the variability of O<sub>2</sub> deficiency at any given site on seasonal as well as inter-annual time scales (Monteiro et al., 2006, 2008).

The region is important for redox nitrogen transformations (Kuypers et al., 2005; Lavik et al., 2009). Surprisingly, however, to our knowledge there have not been any measurements of N<sub>2</sub>O in the water column in this area. On the other hand, CH<sub>4</sub> biogeochemistry in the region has attracted considerable attention largely due to “gas eruptions” (presumably a mixture of CH<sub>4</sub> and hydrogen sulphide (H<sub>2</sub>S)) that usually occur in the late summer (Weeks et al., 2002; Emeis et al., 2004; Brüchert et al., 2006). These eruptions are large enough to be seen by a satellite (Weeks et al., 2002) and the sediment mobilization they cause can occasionally give birth to ephemeral mud islands (Copenhagen, 1953). Several





**Fig. 2.** Time series records of dissolved O<sub>2</sub> (green) and methane (orange) obtained with a mooring deployed off Namibia (Lat 23° S) at 120 m water depth with sensors placed at 85 m. Data are from Monteiro et al. (2006).

sedimentary profiles of CH<sub>4</sub> have been published from the region (water depths 27–109 m – Emeis et al., 2004; Brüchert et al., 2006). High primary production leads to an accumulation of organic carbon and consequently high (anaerobic) respiration in sediments (Brüchert et al., 2006, and references therein). Bacterial reduction of sulphate coupled to degradation of organic matter is intense in the sediment, especially in the upper 10 cm. Sulphate gets fully consumed and methanogenesis occurs at relatively shallow depths (tens of centimeters to a few meters). Accordingly, H<sub>2</sub>S and CH<sub>4</sub> accumulate in porewaters in high concentrations (to ~22 and 8 mM, respectively – Brüchert et al., 2006). Therefore, the source of both gases in the overlying water column in all likelihood is sedimentary even though the relative importance of eruptions/ebullition and diffusion in supplying these gases to the overlying water column is not quite clear. Emeis et al. (2003) suggested that the gases might be released from unconsolidated sediments following physical changes in the sediment or in the overlying water column or even pressure changes arising from rainfall over land that could be transmitted to the sediment through fossil river beds. Obviously such inputs are episodic and even in their absence the water-column build-up of H<sub>2</sub>S can be explained by diffusive fluxes alone (Brüchert et al., 2006, Lavik et al., 2009). This probably applies to CH<sub>4</sub> as well.

Scranton and Farrington (1977) investigated distribution of CH<sub>4</sub> in the water column off Walvis Bay and recorded high values (reaching up to 879 nM) both close to the surface and in near-bottom waters, which were sulphidic over the inner shelf despite little upwelling occurring during the period of the observations (late December 1975–early January 1976). At mid-depth, lower CH<sub>4</sub> concentrations (forming a minimum) probably resulted from the advection of off-shore water. The high concentrations in near-surface waters could arise from inputs from sediments at very shallow depths followed by lateral mixing/offshore advection of

CH<sub>4</sub>-rich waters. However, Scranton and Farrington (1977) opined that some in situ production of CH<sub>4</sub> was probably also occurring in the upper water column. Very recently Brüchert et al. (2009) have provided additional chemically-measured CH<sub>4</sub> data from four stations over the shelf (water depths 27–67 m) that were occupied repeatedly from May 2001 to May 2004. Methane concentrations observed by these authors were even higher – up to 2.9 μM at the surface and 5.2 μM at depth at the shallowest station. To our knowledge, these concentrations are by far the highest reported from any open coastal region.

Results of a high-resolution (hourly) time series, obtained with a mooring having O<sub>2</sub> and CH<sub>4</sub> sensors at 85 m depth, deployed for one year in the same area reveal a tight covariance between the two gases (Monteiro et al., 2006). Methane could not be detected until the O<sub>2</sub> content had declined to 0.2 ml l<sup>-1</sup> in early March 2003 (Fig. 2). The concentrations remained elevated for an extended period thereafter (from May to August), but with brief interruptions associated with increases in the O<sub>2</sub> concentration. The seasonal-scale CH<sub>4</sub> enrichment in bottom waters is modulated by event-scale advection of water as well as by daily-time-scale fluxes of O<sub>2</sub>. Monteiro et al. (2006) proposed that anoxia off Namibia is initially triggered by the advection of equatorial hypoxic waters and then sustained by the respiration of locally produced organic matter.

#### 4.2 Northern Indian Ocean

The northern Indian Ocean comprises two major basins, the Arabian Sea in the northwest and the Bay of Bengal in the northeast. These basins experience very different hydrographic and climatic conditions. The Arabian Sea is a region of negative water balance where evaporation far exceeds precipitation and runoff while the reverse holds true for the Bay of Bengal. Moreover, the Southwest Monsoon winds are also stronger over the Arabian Sea, forcing upwelling along both the western (off Somalia, Yemen and Oman) and eastern (off India) boundaries (Naqvi et al., 2006a). Both the Arabian Sea and the Bay of Bengal experience severe oxygen depletion at mid-depths. While the minimum O<sub>2</sub> concentrations determined by the Winkler method do not differ by more than 0.05 ml l<sup>-1</sup> (~2 μM), this subtle difference is responsible for the prevalence of contrasting redox conditions in the two basins: Unlike the Arabian Sea, a secondary nitrite maximum associated with denitrification is not observed in the Bay of Bengal. Consequently, NO<sub>3</sub><sup>-</sup> concentrations within the core of the OMZ are higher by a factor of ~2 in the Bay of Bengal as compared to the Arabian Sea (Naqvi et al., 2006a). The vertical extent of the OMZ is also smaller in the Bay of Bengal than in the Arabian Sea. Since the northern Indian Ocean is surrounded by land masses on three sides and the O<sub>2</sub> deficiency extends over a wide depth range, the OMZs impinge upon a very large area of the continental margin in this region: as much as two-thirds of the global

continental margin area in contact with bottom waters having O<sub>2</sub> < 0.2 ml l<sup>-1</sup> is found here (Helly and Levin, 2004).

Despite the enormous river runoff into the Bay of Bengal/Andaman Sea and huge consumption of synthetic fertilizers in South Asia, the total flux of dissolved inorganic nitrogen by rivers to the Bay of Bengal is relatively modest (<0.5 Tg N a<sup>-1</sup> – Naqvi et al., 2010a). This is one reason why hypoxic conditions are not known to develop over the inner shelf off the mouths/deltas of major rivers (e.g. Ganges/Brahmaputra and Irrawaddy) unlike, for example, the Gulf of Mexico; the other is that upwelling is very weak and upwelled water does not reach sufficiently close to the surface. By contrast, O<sub>2</sub>-depleted waters ascend to very shallow depths both along the western and eastern boundaries of the Arabian Sea (Naqvi et al., 2006a). There are, however, four important differences between the western and eastern boundary upwelling environments in the Arabian Sea: (1) Upwelling is far more vigorous in the west, which is, in fact, the only major western-boundary upwelling system in the world. Driven by strong southwesterly winds, the strong Ekman flow quickly transports upwelled water offshore to a distance exceeding 1000 km (Naqvi et al., 2006a). (2) Water upwelling in the western Arabian Sea is derived from the south and has relatively high initial O<sub>2</sub> concentration. (3) Unlike the eastern Arabian Sea, there is no freshwater runoff and the upper layer is very weakly stratified in the western Arabian Sea. (4) The continental shelf in the western Arabian Sea is generally narrow which together with vigorous upwelling keeps the residence time of upwelled water over the shelf quite short. The western Arabian Sea, therefore, does not experience the kind of O<sub>2</sub> depletion (suboxic and anoxic conditions in the water column) that distinguishes the western continental shelf of India.


During the Southwest Monsoon the eastern Arabian Sea behaves like a mini EBUE with the equatorward surface flow (as against the poleward flow during the Northeast Monsoon), a poleward undercurrent, and upwelling that is both locally and remotely forced (as against downwelling during the Northeast Monsoon) (Naqvi et al., 2006b, c). However, a thin (<10 m) freshwater lens, formed as a consequence of intense rainfall over the coastal zone, usually prevents the upwelled water from surfacing. The upwelled water is drawn from the undercurrent just off the shelf break that has O<sub>2</sub> content marginally above suboxia. Thus, the combination of sluggish upwelling, low initial O<sub>2</sub> content of the upwelled water, wide shelf, and very strong thermohaline stratification at very shallow depths leads to the formation of the largest natural coastal hypoxic zone (area ~200 000 km<sup>2</sup>) anywhere in the world (Naqvi et al., 2000). It is thus best suited to investigate the effect of coastal hypoxia on the cycling of CH<sub>4</sub> and N<sub>2</sub>O.

#### 4.2.1 Methane

High saturations of CH<sub>4</sub> (124–286‰) in waters upwelled to the surface in the western Arabian Sea have been observed by a number of investigators (Owens et al., 1991; Bange et al., 1998; Upstill-Goddard et al., 1999). These studies focused only on CH<sub>4</sub> distribution at the sea surface. Therefore, even though the bottom waters over the Omani shelf may sometimes become hypoxic during the late Southwest Monsoon (Naqvi et al., 2010b), the contribution of hypoxia in maintaining observed high surface CH<sub>4</sub> concentrations cannot be evaluated. However, as elevated CH<sub>4</sub> concentrations also occur in upwelled waters outside hypoxic zones (e.g. off Mauritania – Kock et al., 2008), at least a part of CH<sub>4</sub> is expected to be produced in the water column (Owens et al., 1991). It may also be noted that surface waters in the Arabian Sea are always phosphate-replete because of large-scale pelagic denitrification in the region (Naqvi, 1987; Morrison et al., 1998; Codispoti et al., 2001). Yet, the upper layer CH<sub>4</sub> maximum is prominently found in the region (Owens et al., 1991), and so its formative mechanism should be different from that proposed by Karl et al. (2008).

Jayakumar et al. (2001) measured CH<sub>4</sub> along a number of coast-perpendicular transects off the central and southwest coast of India during the Southwest Monsoon of 1997 to add to measurements made along a longer transect off Goa during the Spring Intermonsoon of the previous year. Figure 3 shows a typical cross-shelf CH<sub>4</sub> section along with corresponding sections of temperature, salinity, oxygen and nutrients off Goa for the Southwest Monsoon. These sections provide examples of the aforementioned occurrence of strong thermohaline stratification (Fig. 3a, b) and the development of extreme O<sub>2</sub> deficiency within a few meters of the sea-surface (Fig. 3c).

The most striking feature of CH<sub>4</sub> distribution (Fig. 3e) is the sharp onshore-offshore gradient with concentrations decreasing offshore at all depths. Measurements during the Spring Intermonsoon yielded relatively lower and spatially more uniform (3.5–5.5 nM) concentrations (Jayakumar et al., 2001). Higher concentrations during the Southwest Monsoon can arise from two different sources – transport from coastal wetlands and diffusion from underlying sediments. The former source is predominant close to the coast as evident from the association of peak CH<sub>4</sub> values (reaching up to 48 nM, corresponding to over 2500‰ saturation) with the low-salinity cap observed along a closely-sampled shallow transect off Goa (Jayakumar et al., 2001). Even higher CH<sub>4</sub> levels (up to 248 nM, ~13 000‰ saturation at ~15 salinity) were recorded by these authors within the Mandovi Estuary. Thus, over the inner shelf (water depths generally <20 m), CH<sub>4</sub> maximum is located at/near the surface, although concentrations remain elevated even below the pycnocline, which in part may be due to some vertical mixing. At greater water depths, the highest values are found close to the bottom, indicating CH<sub>4</sub> emission from shelf sediments



**Fig. 3.** Distribution of (a) temperature (°C), (b) salinity, (c) O<sub>2</sub> (μM), (d) NO<sub>3</sub><sup>-</sup> (μM), and (e) CH<sub>4</sub> (nM) off Goa (~15.5° N, 73.75° E) during the Southwest Monsoon. Reproduced with permission from Jayakumar et al. (2001).

to the overlying water column. Nevertheless, the observed concentrations are not anomalously high (generally <10 nM) and are comparable to maximal CH<sub>4</sub> concentrations in the upper water column of the open Arabian Sea (Owens et al., 1991; Jayakumar et al., 2001). However, a significant shift in CH<sub>4</sub> distribution is observed with the appearance of H<sub>2</sub>S in bottom waters. This usually happens by late August/early September when nitrate gets fully consumed through denitrification/anammox over the inner- and mid-shelf regions (at water depths of approximately 20–50 m) (Naqvi et al., 2006b, c). The sulphidic bottom waters have been found to contain over 40 nM CH<sub>4</sub> (Gayatree Narvenkar, unpublished data).

The source of CH<sub>4</sub> diffusing from shallow sediment is most likely to be from contemporaneous biogenic production although methane from deeper reservoirs (either petrogenic or biogenic) can also form seeps or plumes. There are reports of gas-charged sediments a few metres below the seafloor off the western coast of India, inferred from acoustic masking. The total amount of CH<sub>4</sub> trapped in such sediments over the inner continental shelf has been estimated as 2.6 Tg

(Karisiddaiah and Veerayya, 1994). However, there has not been any observation of significant CH<sub>4</sub> emission from sediments through bubble ebullition in the region. In any case, the above-mentioned CH<sub>4</sub> inventory does not appear to be sufficient to sustain the observed CH<sub>4</sub> supersaturation in the overlying waters (Jayakumar et al., 2001).

There are few data on CH<sub>4</sub> distribution in sedimentary porewaters off the Indian coast, but the sulphate-CH<sub>4</sub> transition does not appear to be located very close to the surface. Measurements of sedimentary sulphate reduction rates over the inner shelf off Goa yielded surprisingly low values (generally <10 nmol cm<sup>-3</sup> d<sup>-1</sup>). Moreover, porewater sulphide concentrations are also quite low (few micromolar) while sulphate concentrations do not exhibit any significant decrease with depth in the upper 20–30 cm for which data are available (S. W. A. Naqvi and V. Brüchert, unpublished data). Thus, conditions favouring methanogenesis do not seem to occur in the upper few tens of centimeters in sediments. However increased hypoxia in this area, with accompanying increases in carbon flux and lower O<sub>2</sub>, would likely increase the importance of sulphate reduction and CH<sub>4</sub> production within the sediments.

Given the large geographical changes in surface saturation and wind speed, the computed fluxes of CH<sub>4</sub> from the Arabian Sea vary greatly (from ~0 to 64 μmol m<sup>-2</sup> d<sup>-1</sup> – see Naqvi et al., 2005, for a review). The highest emissions are observed during the Southwest Monsoon from waters upwelling in the western Arabian Sea (up to 13.9 μmol m<sup>-2</sup> d<sup>-1</sup> – Owen et al., 1991) and from the inner shelf in the eastern Arabian Sea that is affected by land runoff (up to 64 μmol m<sup>-2</sup> d<sup>-1</sup> – Jayakumar, 1999). Despite these high fluxes, the highest estimate of CH<sub>4</sub> emission from the Arabian Sea as a whole is only 0.1–0.2 Tg a<sup>-1</sup>, which is merely 0.4–0.7% of the total oceanic source and is therefore not very significant (Naqvi et al., 2005).

Methane measurements along three transects over the shelf off Bangladesh in January 1994 by Berner et al. (2003) yielded above-saturation surface concentrations, ranging between 3.17 and 38.3 nM and driving a sea-to-air flux of 0.22–24.9 μmol m<sup>-2</sup> d<sup>-1</sup>. The highest values were observed off the mouth of Ganges/Brahmaputra, reflecting inputs by river water. This is consistent with high CH<sub>4</sub> concentrations of 10.3–59.3 nM reported by Biswas et al. (2003) from the Hooghly Estuary (a distributory of the Ganges). Berner et al. (2003) found hypoxic water (<0.5 ml l<sup>-1</sup>) over the seafloor at depths >60–70 m. Methane concentration in this water reached up to 12.9 nM, so it appears that some enhancement of CH<sub>4</sub> from the sediment underlying hypoxic waters could be occurring over this segment of the shelf. The observed CH<sub>4</sub> concentrations are lower than observed in other regions of river runoff (e.g. in the Gulf of Mexico, see below), in spite of Ganges/Brahmaputra delta containing one of the most extensive mangrove vegetation in the world (the Sundarbans), but then the observations were not made during periods of highest river runoff.




#### 4.2.2 Nitrous oxide

High N<sub>2</sub>O saturations in the Arabian Sea surface waters, averaging 167–186%, were first noticed by Law and Owens (1990) and Naqvi and Noronha (1991). Subsequent studies have led to a large data base on N<sub>2</sub>O distribution in the region that also includes coastal waters off Somalia, Oman and India. An early synthesis of surface measurements by Bange et al. (2001a) shows that the highest concentrations and sea-to-air fluxes occur in the upwelling zones off Oman and India during the Southwest Monsoon. In the western Arabian Sea, surface saturations averaging 230±46% off Oman (in 1994) and up to 330% off Somalia (in 1992) were recorded by Bange et al. (1996), and de Wilde and Helder (1997), respectively. These high values directly arise from high concentration of N<sub>2</sub>O in the upwelling water, which is not suboxic (Naqvi, 1991, 1994) and contains high dissolved N<sub>2</sub>O (Bange et al., 2001b). Observations at 12 stations located over the Omani shelf during September 2004 (S. W. A. Naqvi, unpublished data) yielded surface N<sub>2</sub>O concentrations ranging from 9.7 to 24.7 nM (156–358% saturation), consistent with previous results. However, depth profiles of N<sub>2</sub>O at these stations did not show any anomalously high values, with the maximal subsurface concentration being 48.8 nM. Only at one station was there an indication of mild denitrification occurring in the bottom water with a consequent decrease in N<sub>2</sub>O.

The western continental shelf of India is one of the most interesting and important oceanic sites for N<sub>2</sub>O cycling; arguably it is also the best studied. Distribution of N<sub>2</sub>O over various segments of the coast using data collected on a number of cruises has been investigated (Naqvi et al., 1998, 2000, 2006a, b, c, 2009). In addition, measurements of N<sub>2</sub>O have also been made since 1997 during monthly/fortnightly trips to a coastal quasi-time series station (the Candolim Time Series – CaTS) located off Goa (Lat. 15°31' N, Long. 73°39' E) at a water depth of ~26 m. The monthly/fortnightly-averaged records showing annual cycles of key oceanographic parameters are shown in Fig. 4.

During the Northeast Monsoon and Spring Intermonsoon periods, when the water column over the Indian shelf is well oxygenated (Fig. 4c), N<sub>2</sub>O concentrations do not exceed 10 nM (Fig. 4f). Nitrous oxide begins to accumulate as O<sub>2</sub> is consumed in the subsurface layer after the onset of Southwest Monsoon upwelling (usually in June). The concentrations continue to rise even after the system becomes suboxic. In fact, the greatest build-up of N<sub>2</sub>O coincides with the rapid decline in NO<sub>3</sub><sup>-</sup> (Fig. 4d) and accumulation of NO<sub>2</sub><sup>-</sup> (Fig. 4e), pointing to N<sub>2</sub>O production through denitrification. This is in sharp contrast with the pattern observed in open-ocean OMZs, where, as mentioned above, the secondary nitrite maximum is invariably characterized by a minimum in N<sub>2</sub>O. However, the concentrations decrease rapidly once the environment becomes sulphate reducing (Fig. 4h), which indicates that the observed high concentrations cannot be due



**Fig. 4.** Monthly-/fortnightly-averaged records showing annual cycle of (a) temperature, (b) salinity, (c) oxygen, (d–g) inorganic nitrogen species, and (h) hydrogen sulphide at the Candolim Time Series (CaTS) site (15.52° N, 73.65° E) based on observations from 1997 to 2006. Modified from Naqvi et al. (2009).

to inhibition of N<sub>2</sub>O reductase activity by H<sub>2</sub>S (Senga et al., 2006).

Observations along cross-shelf transects also show a similar pattern of N<sub>2</sub>O variability (Naqvi et al., 2000, 2006b, c, 2009). During the Northeast Monsoon N<sub>2</sub>O concentrations are low and less variable (<25 nM to a depth of 100 m). But the onset and intensification of O<sub>2</sub> deficiency through the summer (Southwest Monsoon) brings about dramatic changes in the N<sub>2</sub>O field. By late summer/early autumn one can identify three zones with different redox conditions prevailing in bottom waters north of about 12° N latitude: hypoxia over the outer shelf (offshore of 50/60-m water depth), suboxia over the mid shelf (between 50/60-m and 20/30-m water depths), and anoxia over the inner shelf (between 20/30-m and 10-m water depths). Both the highest and the lowest N<sub>2</sub>O concentrations are associated with reducing environments. The highest recorded value (765 nM off Goa – Naqvi et al., 2006c), for example, came from a sample that had lost most of its NO<sub>3</sub><sup>-</sup> through denitrification. In addition to the association of high N<sub>2</sub>O and NO<sub>2</sub><sup>-</sup> (up to 16 μM)

values, accumulation of N<sub>2</sub>O during denitrification has also been demonstrated by Naqvi et al. (2000) through incubation in air-tight bags of water samples that were initially close to but not quite suboxic (O<sub>2</sub> ~ 15 µM). It was suggested by these authors that frequent aeration of the water through turbulence could suppress the activity of N<sub>2</sub>O reductase allowing transient production of N<sub>2</sub>O. In regions where, and during periods when, subsurface O<sub>2</sub>-deficiency is not severe enough to allow the onset of suboxic conditions, as happens off the southwest coast of India (generally south of ~12° N latitude), abnormally high (>100 nM) N<sub>2</sub>O concentrations have not been recorded (Naqvi et al., 2006a).

Maximal surface concentration observed over the Indian shelf, which is also the highest reported from the oceanic surface waters, is 436 nM (corresponding to 8250% saturation), with the average exceeding 37 nM (Naqvi et al., 2006b). The highest computed flux is 3243 µmol m<sup>-2</sup> d<sup>-1</sup>, with the average ranging from 39 to 264 µmol m<sup>-2</sup> d<sup>-1</sup> depending upon the model of air-sea exchange chosen and the wind speed (5–10 m s<sup>-1</sup> – Naqvi et al., 2006b). The total emission of N<sub>2</sub>O from the Indian shelf is thus computed to be 0.05–0.38 Tg N<sub>2</sub>O for the upwelling season. For comparison, the total annual efflux of N<sub>2</sub>O from the Arabian Sea as a whole (that did not take into account the abnormally high values from the Indian coast) was estimated to range between 0.33 and 0.70 Tg by Bange et al. (2001a). It has been suggested that the natural low-O<sub>2</sub> system off India has intensified in recent years most likely because of enhanced nitrogen loading through runoff and atmospheric deposition (Naqvi et al., 2000, 2006b, 2009). Because N<sub>2</sub>O data from this region go back only to 1997, it is not clear to what extent has this intensification affected N<sub>2</sub>O cycling. However, given the observation that the highest N<sub>2</sub>O concentration is observed when O<sub>2</sub> deficiency is the most intense, we speculate that an increase in production is likely to have occurred relative to earlier pristine conditions.


Unlike the Arabian Sea, the OMZ of the Bay of Bengal is just short of being suboxic in that a secondary nitrite maximum generally does not occur in the region. Consequently, vertical profiles of N<sub>2</sub>O show a single broad maximum within the OMZ (Naqvi et al., 1994). Surface saturations (89–214%) and atmospheric fluxes (–0.10 to 10.67 µmol m<sup>-2</sup> d<sup>-1</sup>) from the region are, therefore, much lower. Also, although some upwelling does occur along the Indian east coast during the Southwest Monsoon, the shelf is narrow and the low-salinity layer formed as a result of enormous freshwater inputs to the Bay of Bengal through rainfall and river runoff is several tens of meters thick. As a result the upwelled water does not reach very shallow depths (Naqvi et al., 2006a), and maximal N<sub>2</sub>O concentrations over the shelf are a few tens of nanomolar (Naqvi et al., 1994, 2006a).

### 4.3 Eastern North Pacific Ocean

The eastern tropical/subtropical North Pacific experiences conditions that are typical of an oceanic eastern boundary: the surface current (the California Current) flows equatorward while the northwesterly winds drive intense upwelling along the west coast of the United States and the northwest coast of Mexico. An extensive OMZ with O<sub>2</sub> < 0.2 ml l<sup>-1</sup> (9 µM) extends ~1500 km offshore from the Mexican coast (Deuser, 1975). The suboxic zone, confined to the tropical region, is among the best investigated for redox nitrogen cycling (Cline and Richards, 1972; Cline and Kaplan, 1975; Ward et al., 2008). Low-O<sub>2</sub> concentrations also extend at mid-depths quite far north off the west coast of United States (especially in the silled basins of California Borderland), although the water column does not seem to be reducing (e.g. Sigman et al., 2003). As already mentioned, large decreases in subsurface O<sub>2</sub> concentrations in the California Current System have been recorded recently. These are believed to result from advection of O<sub>2</sub> depleted water over the shelf (Grantham et al., 2004; Bograd et al., 2008; Chan et al., 2008; Connolly et al., 2010).

#### 4.3.1 Methane


The open-ocean suboxic zone of the eastern tropical North Pacific (ETNP) contains the largest pool of CH<sub>4</sub> in the open ocean, although the computed sea-to-air flux from this region (0.77–3.0 µmol m<sup>-2</sup> d<sup>-1</sup>) is quite modest (Sansone et al., 2001). Vertical CH<sub>4</sub> profiles obtained by Burke et al. (1983) at several stations located parallel to but away from the Mexican coast, showed a ubiquitous shallow (50–150 m) CH<sub>4</sub> maximum outside the zone of suboxia with concentrations reaching up to 6.5 nM. This maximum extended much deeper (to at least 400 m, which was the maximum depth of sampling) at stations that were located within the suboxic zone. This was attributed to in-situ production of CH<sub>4</sub> through microbial activity associated with suspended particles recycled by zooplankton grazing. It may be noted that a secondary particle (turbidity) maximum was also found within suboxic waters. A subsequent study by Sansone et al. (2001) in the ETNP along a transect extending offshore from the Mexican coast also found relatively higher CH<sub>4</sub> concentrations in the suboxic layer (Fig. 5c). Methane levels were particularly elevated (28 nM at 350 m) at the station located closest to the coast. These authors also measured the carbon isotopic composition of CH<sub>4</sub> and concluded that while CH<sub>4</sub> in the upper half of the pool was being produced in situ during decomposition of sinking organic matter, that in the lower half of the pool was derived from the continental margin. In a more recent study, Sansone et al. (2004) focused on CH<sub>4</sub> cycling along the western Mexican continental margin in and around the Gulf of California (Fig. 6c, d). Their sampling stations were positioned both in the coastal basins and over the open margins. Those stations where water depth exceeded the



**Fig. 5.** Vertical profiles of dissolved O<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub> concentrations, δ<sup>15</sup>N of N<sub>2</sub>O and Site Preference (SP) of <sup>15</sup>N in N<sub>2</sub>O at a station located within the open-ocean suboxic zone of the eastern tropical North Pacific (16° N, 107° W). Data are from Sansone et al. (2001) for CH<sub>4</sub> and from Yamagishi et al. (2007) for N<sub>2</sub>O.

lower boundary of the OMZ had lower CH<sub>4</sub> concentrations than the station where the OMZ impinged upon the seafloor, consistent with higher CH<sub>4</sub> supply to and/or lower oxidative loss within the OMZ. The highest build-up of CH<sub>4</sub> (up to 78 nM) was found to occur in bottom waters of silled basins (Fig. 6d), presumably through diffusion from the seafloor. Apparently this did not affect CH<sub>4</sub> distribution outside the basins, perhaps because of water column CH<sub>4</sub> oxidation. Diffusive fluxes from the sediment (0.24–5.5 μmol m<sup>-2</sup> d<sup>-1</sup>) computed from porewater CH<sub>4</sub> gradients were similar to the sea-to-air fluxes (0.5–5.9 μmol m<sup>-2</sup> d<sup>-1</sup>), and since oxidative loss must be occurring in the water column, a source of CH<sub>4</sub> in the water column is also required to sustain the sea-to-air flux.

Several studies have been conducted on CH<sub>4</sub> cycling in coastal waters further north in the California Current System (e.g. Ward, 1992; Cynar and Yayanos, 1992, 1993; Tilbrook and Karl, 1995; Kessler et al., 2008). The usual trend of CH<sub>4</sub> supersaturation in surface waters with a subsurface maximum has been reported but no major anomalies have been seen that could be attributed to hypoxia. There seems to be general agreement that inputs from the continental margin are important in maintaining water column CH<sub>4</sub> maxima, particularly because there is significant hydrocarbon seepage in the region. For example, Cynar and Yayanos (1992) observed values as high as 1416 nM between Point Conception and Santa Barbara, which could only be produced by CH<sub>4</sub> emission from hydrocarbon seeps. Radiocarbon measurements confirmed the importance of seep-derived CH<sub>4</sub> in the Santa Barbara Basin (Kessler et al., 2008). Due to high CH<sub>4</sub> concentration in subsurface waters, upwelling enhances supersaturation in surface waters (Rehder et al., 2002).



**Fig. 6.** (a–c) Vertical profiles of dissolved O<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub> concentrations, δ<sup>15</sup>N of N<sub>2</sub>O and SP of <sup>15</sup>N in N<sub>2</sub>O at an open margin station in the Gulf of California (26.3° N, 110.13° W); (d) Vertical profiles of dissolved O<sub>2</sub> and CH<sub>4</sub> at a silled-basin station (sill depth 50 m) in the same region (21.3° N, 105.9° W). Data are from Sansone et al. (2001, 2004) for CH<sub>4</sub> and from Yamagishi et al. (2007) for N<sub>2</sub>O.

#### 4.3.2 Nitrous oxide

Early studies of N<sub>2</sub>O in the ETNP were undertaken by Cohen and Gordon (1978) and by Pierotti and Rasmussen (1980). Surface waters were found to be generally supersaturated with respect to atmospheric N<sub>2</sub>O (on an average by 111% and 123% in the two studies), driving a flux of N<sub>2</sub>O from the ocean to the atmosphere. Vertical N<sub>2</sub>O profiles, more comprehensively examined by Cohen and Gordon (1978), revealed the existence of double maxima in the region affected by denitrification. This was attributed to consumption of N<sub>2</sub>O by denitrifiers within the secondary nitrite maximum and its production at the boundaries by nitrifiers, a view largely supported by subsequent measurements including dual isotopic composition of N<sub>2</sub>O (Yoshinari et al., 1997; Yamagishi et al., 2007). Yamagishi et al. (2007) presented data from two stations – one each located within the heart of the suboxic zone of the ETNP (Fig. 5a, b) and the central Gulf of California (Fig. 6a, b) – on N<sub>2</sub>O concentration and its isotopic composition. The latter also included isotopomeric analysis (i.e. the location of <sup>15</sup>N in linear N<sub>2</sub>O molecule) which provides information on the mechanisms of N<sub>2</sub>O production. In the ETNP, the highest N<sub>2</sub>O concentration (~87 nM, Fig. 5a) and the lowest isotopic values (e.g. δ<sup>15</sup>N<sub>bulk</sub>=3.6‰ vs. air, Fig. 5b) occurred at the lower oxycline (depth 65 m, O<sub>2</sub>=32 μM) whereas within the core of the secondary nitrite maximum where the N<sub>2</sub>O concentration fell well below saturation, N<sub>2</sub>O was isotopically very heavy (δ<sup>15</sup>N<sub>bulk</sub>=22.7‰ vs. air at 400 m). In the Gulf of California, the double maximum characteristic of suboxic zones was not observed (note that a secondary nitrite maximum was also not reported at this station). Between 300 and 970 m water depths, where O<sub>2</sub> concentration varied from 0.6 to 3.1 μM, N<sub>2</sub>O concentration was in the range 41.6–84.4 nM (Fig. 6a). Interestingly, isotopic values (Fig. 6b) varied considerably

over this depth range (e.g.  $\delta^{15}\text{N}^{\text{bulk}}$  was between 1.76 and 1.94‰ at 705–805 m where N<sub>2</sub>O reached peak values, as compared to 10.97‰ at 300 m). The conventional interpretation of these data would be that greater production of N<sub>2</sub>O occurred through nitrification at 700 m as compared to 300 m. However, isotopomeric measurements suggest an opposite trend according to which most of the production of N<sub>2</sub>O in deeper waters of the Gulf of California should be through denitrification (Yamagishi et al., 2007).

Another example of N<sub>2</sub>O behaviour in low-O<sub>2</sub> waters in the eastern North Pacific comes from the work of Codispoti et al. (1992) who generated data on N<sub>2</sub>O concentration along a section extending about 250 km offshore from Monterey Bay/Canyon, California. The isotopic composition of N<sub>2</sub>O was also determined at a station toward the shoreward end of the transect. The water was hypoxic below ~300 m, with O<sub>2</sub> < 12.5 µM between 600–800 m where N<sub>2</sub>O concentrations exceeded 47.5 nM. The  $\delta^{15}\text{N}$  and  $\delta^{18}\text{O}$  values mostly hovered around those in the troposphere ( $\delta^{15}\text{N}=7.0\text{‰}$  vs. air and  $\delta^{18}\text{O}=44.2\text{‰}$  vs. SMOW), but consistent maxima in both properties (e.g. up to 9‰ for  $\delta^{15}\text{N}$ ) were found within the layer of minimum O<sub>2</sub> and maximum N<sub>2</sub>O. This is at variance with the above-mentioned trend observed in the Gulf of California, and shows how minor changes in O<sub>2</sub> concentration can result in dramatic changes in N<sub>2</sub>O concentration and its isotopic composition (and consequently the production mechanisms).

Estimates of sea-to-air fluxes from the ETNP vary over a wide range. Cohen and Gordon (1978) reported an average efflux of  $2.75\text{ }\mu\text{mol m}^{-2}\text{ d}^{-1}$  (range  $-0.79$ – $5.5\text{ }\mu\text{mol m}^{-2}\text{ d}^{-1}$ ), which is substantially lower than the corresponding estimate ( $10.6\text{ }\mu\text{mol m}^{-2}\text{ d}^{-1}$ ) of Pierotti and Rasmussen (1980). Using the same data set (of Pierotti and Rasmussen), however, Cohen and Gordon (1978) computed an average efflux of  $6.68\text{ }\mu\text{mol m}^{-2}\text{ d}^{-1}$  between 4 and 22° N latitudes. Accordingly, the estimate for total N<sub>2</sub>O emission from the ETNP by these authors should also be scaled down to  $0.75\text{ Tg N}_2\text{O a}^{-1}$ . For comparison, the average value of Cohen and Gordon (1978) when applied to the same area yields an emission estimate of  $\sim 0.3\text{ Tg N}_2\text{O a}^{-1}$ .

#### 4.4 Eastern South Pacific Ocean

Circulation along the western boundary of South America, described in detail by Strub et al. (1998), consists of the surface Humboldt Current carrying low-salinity and well-oxygenated Subantarctic Water (SAAW) equatorward and, the subsurface Peru-Chile Undercurrent transporting high-nutrient, high-salinity, and low-oxygen Equatorial Subsurface Water (ESSW) poleward. The southerly and south-westerly winds cause Ekman transport away from the coast, and depending on the wind intensity, bottom topography and other oceanographic conditions associated principally with remote forcing, e.g. El Niño – Southern Oscillation (ENSO) (Shaffer et al., 1999), the ESSW upwells along the Peruvian-

Chilean coast, bringing cold, nutrient-rich, oxygen-poor waters to the surface. The consequent high biological production sustains one of the world's most intense OMZs that extends far offshore into the open ocean from very shallow depths. The intensity and thickness of the OMZ, which is associated with the ESSW, decreases with the feature occurring progressively deeper toward the south. Thus, the most intense O<sub>2</sub>-deficiency occurs near Peru, where episodes of complete denitrification and associated sulphidic conditions in the water column are recorded in the literature (e.g. Dugdale et al., 1977; Codispoti et al., 1986).


##### 4.4.1 Methane

The area off central Chile experiences upwelling and bottom-water O<sub>2</sub> depletion on a seasonal basis. This area has several topographic and oceanographic characteristics that set it apart from the other EBUEs. The continental shelf is quite wide here. In the Austral spring and summer, intensified winds from the south and southwest make the ESSW ascend over this shelf (Sobarzo and Djurfeldt, 2004). As a result, the photic zone gets greatly enriched with nutrients, leading to high, but very variable, primary production ( $1$ – $19\text{ g C m}^{-2}\text{ d}^{-1}$ ; Daneri et al., 2000) and consequently intense respiration of organic matter in bottom waters.

Methane measurements have been made since 2007 at a time-series station located over the continental shelf off Concepción, central Chile, at a water depth of ~90 m (Fig. 7a). The observed concentrations vary from 3 to 70 nM. Methane profiles usually show a significant increase from the surface (3–19 nM, 123–750% saturation) to the bottom (36–70 nM). Methane fluxes across the air-sea interface in this area are always directed from the ocean to the atmosphere, ranging between  $0.05$  and  $59.5\text{ }\mu\text{mol m}^{-2}\text{ d}^{-1}$  (mean  $25.6\text{ }\mu\text{mol m}^{-2}\text{ d}^{-1}$  – L. Farias, unpublished data). The emissions are maximal during upwelling periods. These values are higher than the published fluxes from the Arabian Sea (Owens et al., 1991; Bange et al., 1998; Upstill-Goddard et al., 1999; Jayakumar et al., 2001).

Temporal variability of vertical CH<sub>4</sub> distribution (Fig. 7a) is characterized by marked seasonality, with the bottom water concentrations peaking during the upwelling-favorable periods in synchrony with a drop in O<sub>2</sub> levels. This is in agreement with the above-mentioned observations off Namibia, suggesting greater CH<sub>4</sub> diffusion from the sediment and/or smaller consumption (oxidative loss) in O<sub>2</sub>-depleted bottom waters. However, there are times when subsurface water has lower CH<sub>4</sub> concentration than surface water. These may indicate either advective input of lower CH<sub>4</sub> offshore waters or a dynamic balance between production/supply and loss terms. Incubation of samples with labelled tracer yielded negative net CH<sub>4</sub> cycling rates, thereby providing evidence for dominance of methanotrophy (net CH<sub>4</sub> consumption) in the water column (Farias et al., 2009b).





**Fig. 7.** Temporal variability of (a) CH<sub>4</sub>, (b) O<sub>2</sub>, (c) N<sub>2</sub>O; and (d) NO<sub>2</sub><sup>-</sup> at the COPAS time series station located over the shelf off central Chile at a water depth of ~90 m. Data are from Farias et al. (2009b) for CH<sub>4</sub> and from Cornejo et al. (2007) for other parameters.

#### 4.4.2 Nitrous oxide

##### Off Peru and northern Chile (10°–23° S)

The area off northern Chile and Peru experiences quasi-permanent coastal upwelling and contains one of the shallowest and most intense oceanic OMZs. A sharp oxycline, developed as the combined consequence of advection of low-

O<sub>2</sub> ESSW and break-down of copious amounts of locally-produced organic matter, is thus located at relatively shallow depths, often within the photic zone (Paulmier et al., 2006). Below the oxycline to a depth of ~400 m, the O<sub>2</sub> concentration is close to zero (Revsbech et al., 2009). The secondary nitrite maximum formed due to dissimilatory reduction of NO<sub>3</sub><sup>-</sup> occurs within the OMZ (Codispoti et al., 1986).

As in other suboxic zones, vertical N<sub>2</sub>O distribution in this region shows marked extrema. The concentrations are the highest (reaching up to 86 nM) within the oxycline, accounting for as much as 41–68% of the  $\Delta$ N<sub>2</sub>O pool. The minimum (10–25 nM) occurs within the secondary nitrite maximum, located between 100 and 300 m, beneath which a secondary N<sub>2</sub>O maximum is found as in other areas (Fariás et al., 2007; also see Codispoti et al., 1992). The  $\delta^{15}\text{N}$ -N<sub>2</sub>O increases from 8.57‰ (vs. air) within the oxycline (50 m depth) to 14.87‰ (100 m depth) within the secondary nitrite maximum, the range being somewhat narrower than in the ETNP (3.6 to 22.7‰ – Yamagishi et al., 2007) and the Arabian Sea (–2 to 81‰ – Naqvi et al., 2006c). Based on incubations of samples collected from the oxycline, Farias et al. (2009a) estimated N<sub>2</sub>O production rates through denitrification (NO<sub>2</sub><sup>–</sup> reduction to N<sub>2</sub>O) ranging from 2.25 to 50.05 nmol l<sup>–1</sup> d<sup>–1</sup>. Nitrous oxide should also be produced by aerobic ammonium oxidation within the oxycline given that up to 42% of NH<sub>4</sub><sup>+</sup> produced in this layer is oxidized by nitrifiers off northern Chile (Molina et al., 2005; Molina and Farias, 2009). However, the relative importance of these processes in N<sub>2</sub>O production is difficult to quantify. Farias et al. (2009a) also measured N<sub>2</sub>O consumption rates (i.e. its reduction to N<sub>2</sub>) of 2.73–70.8 nmol l<sup>–1</sup> d<sup>–1</sup>, with this process becoming progressively more important toward the core of the OMZ. Thus, N<sub>2</sub>O turnover should be quite rapid.

Coastal waters off Peru exhibit some of the highest concentrations of N<sub>2</sub>O observed in the ocean at or close to sea surface. Based on measurements made in March 1978, Elkins (1978, as cited by Codispoti et al., 1992) observed an average saturation of 632% in surface waters in the region south of about 12° S latitude. Codispoti et al. (1992) themselves found an average saturation of 450% with the maximum surface concentration of 173 nM. They noted that their observations were made during a period of weak upwelling (in February–March 1985) and that higher concentrations would be expected during more vigorous upwelling. Significantly, the highest N<sub>2</sub>O concentration in subsurface waters (175–195 nM) are from shallow, high-NO<sub>2</sub><sup>–</sup> (most probably suboxic) waters (see Fig. 1B in Codispoti et al., 1992). This is very similar to observations over the western Indian shelf and would support the view that such high concentrations represent transient accumulation of N<sub>2</sub>O during denitrification at shallow depths. As in the case of the Arabian Sea, the high surface concentrations feed large efflux of N<sub>2</sub>O to the atmosphere (12.7–30.7  $\mu\text{mol m}^{-2} \text{d}^{-1}$  – Farias et al., 2009a). It may be noted that negative sea-to-air fluxes, implying net N<sub>2</sub>O flux into the ocean, have also been reported from the region (Table 2b) (Pierotti and Rasmussen, 1980). Codispoti et al. (1992) computed the total annual emission from their area of observations (250 000 km<sup>2</sup>) to be 0.16 Tg N<sub>2</sub>O. This is much smaller than the total annual emission (0.8 Tg N<sub>2</sub>O) estimated by Pierotti and Rasmussen (1980) for the entire O<sub>2</sub>-deficient region of the ETSP. However, the latter is prob-

ably an overestimate in view of the systematically high N<sub>2</sub>O concentrations measured by these authors in the atmosphere for which corrections have been made in the data included in Table 2b.

### Off Central Chile (30°–40° S)

Regular monitoring of physical and biogeochemical conditions at the above-mentioned COPAS time series site included N<sub>2</sub>O since its inception in 2002. The time series records show the recurrence of hypoxia/suboxia in the water column every summer (Fig. 7b). The onset of O<sub>2</sub> deficiency affects N-biogeochemistry both in the water column and sediments, and as in other shallow systems, a close coupling between pelagic and benthic processes is to be expected (Fariás et al., 2004).

As anticipated, N<sub>2</sub>O saturation in surface water is the highest (up to 1372%) during spring-summer upwelling period, whereas minimal values (sometimes below saturation) occur during winter. Peak accumulation of N<sub>2</sub>O is observed mostly at mid-depth (within the oxycline) and sometimes in suboxic bottom waters, with considerable inter-annual variability. The highest concentration recorded is 245 nM. Such “hotspots” of N<sub>2</sub>O production (Fig. 7c) occur during periods of most intense O<sub>2</sub> deficiency (Fig. 7b) and accumulation of NO<sub>2</sub><sup>–</sup> in bottom waters (Fig. 7d) (Cornejo et al., 2007). The association of high N<sub>2</sub>O values with varying O<sub>2</sub> levels points to production through nitrification as well as denitrification. Results of measurements of carbon assimilation in the dark, with and without the addition of allylthiourea, an inhibitor of monooxygenase enzymes of aerobic ammonium oxidation, and the natural isotope abundance in particulate organic carbon ( $\delta^{13}\text{C}$ -POC) showed that chemolithoautotrophs and specifically aerobic ammonium oxidizers are active in the region, especially during the upwelling period, facilitating elevated production of N<sub>2</sub>O (Farias et al., 2009b).

The onset of hypoxic/suboxic conditions affects the sediment-water exchange of dissolved nitrogen species (Farias and Cornejo, 2007). Conditions in bottom water are influenced by benthic organic remineralization, which consumes O<sub>2</sub> as well as other electron acceptors such as NO<sub>3</sub><sup>–</sup>. Benthic fluxes of NO<sub>3</sub><sup>–</sup> (2.62–5.08 mmol m<sup>–2</sup> d<sup>–1</sup>) and N<sub>2</sub>O (4.46–5.53  $\mu\text{mol m}^{-2} \text{d}^{-1}$ ) are always directed into the sediments when hypoxic/suboxic conditions prevail in overlying waters. Nitrous oxide consumption in sediments occurs even during oxic conditions, but the rate decreases by a factor of 2.

Compared to the northern coast of Chile (off Iquique), N<sub>2</sub>O efflux off central Chile exhibits marked seasonality (Cornejo et al., 2007). The upwelling period is characterized by N<sub>2</sub>O emission from the ocean to the atmosphere (up to 195  $\mu\text{mol m}^{-2} \text{d}^{-1}$ ), while low, even negative values of sea-to-air flux (to –9.8  $\mu\text{mol m}^{-2} \text{d}^{-1}$ ) are obtained for the non-upwelling period (May–July), with the annual average exceeding 10  $\mu\text{mol m}^{-2} \text{d}^{-1}$  (Cornejo et al., 2007; L. Farias, unpublished). The zonal distribution of N<sub>2</sub>O flux across



**Table 2a.** Concentrations and sea-to-air fluxes of CH<sub>4</sub> from major hypoxic/suboxic/anoxic zones.

Region	Surface/mixed layer concentration (nM)	Maximum subsurface (depth) concentration (nM)	Sea to air flux ( $\mu\text{mol m}^{-2} \text{d}^{-1}$ )	References
<b>Normoxic Open Ocean</b>	Slightly supersaturated with respect to atm. CH <sub>4</sub> (e.g. $\sim 2$ nM in the tropics)	Maximum at the base of the mixed layer (e.g. $\sim 3$ nM in the tropics)	Variable, but mean $< 0.5 \mu\text{mol m}^{-2} \text{d}^{-1}$	Bates et al. (1996); Reeburgh (2007); Rhee et al. (2009)
<b>Deep Hypoxic/Suboxic Systems</b>				
Eastern Tropical North Pacific	1.8–4.1	6.5 (140 m)	Not given (N.G.)	Burke et al. (1983)
	2.3–4	28 (350 m)	0.77–3	Sansone et al. (2001)
Eastern Tropical South Pacific	2.1–4	12.8 (150 m)	$\sim 0$ –4.3	Kelley and Jeffrey (2002)
Arabian Sea	$< 2$ –5.3	10.1 (35 m), 8.2 (200 m)	$\sim 0$ –13.9	Owens et al. (1991); Jayakumar et al. (2001); Naqvi et al. (2005)
Bay of Bengal ( $> 10^\circ \text{N}$ )	1.7–4.1	6.5 (75 m)	–0.9–4.7	Berner et al. (2003)
<b>Naturally Hypoxic Shelf Systems (including deeper basins)</b>				
Eastern Tropical North Pacific				
Mexican margin (open)	3.2–5.7	46 (340 m)	0.5–2.7	Sansone et al. (2004)
Mexican margin (Gulf of California basins)	3–4.5	25–78 (400–530 m)	0.5–5.9	Sansone et al. (2004)
California Borderland basins	2.9–10.5 <sup>a</sup>	$\sim 50$ (50 m, 550 m) <sup>a</sup>	0.1– $> 147^a$	Ward (1992); Cynar and Yayanos (1992, 1993); Kessler et al. (2008)
Eastern Tropical South Pacific (Off central Chile)	3.1–29.2	79.9 (90 m)	0.05–59.5	Farias et al. (2009b); L. Farias, unpublished
Eastern Tropical North Atlantic (Off Mauritania)	2–5	N.G.	0.5–0.8	Kock et al. (2008)
Eastern Tropical South Atlantic (off Namibia)	2.2–2870	5160 (18 m) 200–2000 (Benthic Boundary Layer)	3–752 N.G.	Scranton and Farrington (1977) Monteiro et al. (2006) Brüchert et al. (2009)
Eastern Arabian Sea (Indian shelf)	2.6–47.6	44 (24 m)	0.58–63.7	Jayakumar (1999); G. Narvenkar, unpublished
Northern Bay of Bengal	3.2–38.3	12.9 (78 m)	0.22–24.9	Berner et al. (2003)
<b>Anthropogenically-produced Coastal Hypoxic Systems</b>				
Gulf of Mexico	$< 3$ –750	708 (15 m)	N.G.	Kelley (2003)
Changjiang (Yangtze River) Estuary and East China Sea	2.8–49	85.9 (5 m)	0.02–440	Zhang et al. (2008b)
<b>Land-locked Anoxic Basins</b>				
<b>Baltic Sea</b>				
Eckernförde Bay (Pockmark site)	2–106	441 (25 m)	–0.4–413	Bussmann and Suess (1998)
Eckernförde Bay (Boknis Eck Time Series Station)	7–42	235 (20 m)	6–15	Bange et al. (2010)
Kiel Fjord	80–1000	8000 (10 m)	N.G.	Schmaljohann (1996)
Baltic Proper	4–695	N.G.	10–1200	Bange et al. (1994)
Baltic Proper	4–22	403 (depth N.G.)	N.G.	Dzyuban et al. (1999)
Mariager Fjord	500–900	30 000 (30 m)	240–4500	Fenchel et al. (1995)
<b>Black Sea</b>				
Northwest, largely shelf	4–255	200 (50 m)	32–470	Amouroux et al. (2002)
Deep basin	10	13 050 (1650 m)	26.6–50	Amouroux et al. (2002); Reeburgh et al. (1991); Kessler et al. (2006a)
Cariaco Basin	$\sim 2$	16 780 (1370 m)	0.23	Ward et al. (1987); Kessler et al. (2006b)
Saanich Inlet	23.4–50	1609 (175 m)	2.3–57	Bullister et al. (1982); Lilley et al. (1982); Ward et al. (1989)

<sup>a</sup> Except stations located very close to coast sampled by Cynar and Yayanos (1992, 1993) where surface and subsurface values reached up to  $\sim 1200$  nM and 1400 nM, respectively, with sea-to-air fluxes as high as  $4350 \mu\text{mol m}^{-2} \text{d}^{-1}$ .

**Table 2b.** Concentrations and sea-to-air fluxes of N<sub>2</sub>O from major hypoxic/suboxic/anoxic zones.

Region	Surface/mixed layer concentration (nM)	Maximum subsurface (depth) concentration (nM)	Sea to air flux ( $\mu\text{mol m}^{-2} \text{d}^{-1}$ )	References
<b>Normoxic Open Ocean</b>	Slightly supersaturated with respect to atm. N <sub>2</sub> O (e.g. ~5–6 nM in the tropics)	Coincides with the oxygen minimum (e.g. ~30 nM at ~500 m in the equatorial Atlantic)	Variable but mean $<1 \mu\text{mol m}^{-2} \text{d}^{-1}$	Butler et al. (1989); Nevison et al. (1995); Rhee et al. (2009)
<b>Deep Hypoxic/Suboxic Systems</b>				
Eastern Tropical North Pacific	4.9–11.1	86.9 (65 m)	–5.6–15.9	Cohen and Gordon (1978); Pierotti and Rasmussen (1980); Yamagishi et al. (1997)
Gulf of California	12.5	84.4 (705 m)	N.G.	Yamagishi et al. (1997)
Eastern Tropical South Pacific (Off Peru and northern Chile)	6.1–37.8	86 (30 m)	–1.1–30.7	Pierotti and Rasmussen (1980); Farias et al. (2009b)
Arabian Sea	5.2–16.7	85.6 (800 m)	~0–470	Naqvi et al. (2005)
Bay of Bengal	4.95–7.6	79.6 (205 m)	0.6	Naqvi et al. (1994)
<b>Naturally Hypoxic Shelf Systems</b>				
Eastern Tropical South Pacific				
Off Peru	< 11.4–172.7	195.5	~39	Codispoti et al. (1992)
Off central Chile	6.7–58.9	245 (30 m)	–9.8–195	Cornejo et al. (2007)
	5.1–30.1	206 (40 m)	–7.7–42.9	Farias et al. (2009a); L. Farias, unpublished
Arabian Sea				
Omani shelf	9.7–24.7	48.8 (50 m)	N.G.	S. W. A. Naqvi, unpublished
Indian shelf	5.3–436	765 (19 m)	–1.2–3243	Naqvi et al. (2006b)
Western Bay of Bengal	4.9–12.0	42.0 (120 m)	–0.1–10.7	Naqvi et al. (1994)
<b>Anthropogenically-produced Coastal Hypoxic Systems</b>				
Gulf of Mexico	5–7	47	0.6–11.7	Walker et al. (2010)
Changjiang (Yangtze River) Estuary and East China Sea	5.8–37.2	62.6 (20 m)	–0.7–97.5	Zhang et al. (2008b)
Chesapeake Bay	6.6–9.8	Highest value at the surface	N.G.	Elkins et al. (1978)
Tokyo Bay	8.8–139	Highest value at the surface	1.51–153	Hashimoto et al. (1999)
<b>Land-locked Anoxic Basins</b>				
Baltic Sea				
Baltic Proper	14–20	1523 (110 m)	5–11	Rönnner (1983)
Gotland Basin	13	63 (90 m)	N.G.	Brettar and Rheinheimer (1991)
Baltic Proper	10–12	31 (90 m)	N.G.	Walter et al. (2006)
Eckernförde Bay (Boknis Eck Time Series Station)	10–17	19 (25 m)	N.G.	Schweiger (2006)
Black Sea				
Northwest shelf	6.5–8	N.G.	1.6–4.4	Amouroux et al. (2002)
Deep basin	7.5–10.2	14.4 (70 m)	3.1–5.2	Amouroux et al. (2002); Butler and Elkins (1991); Westley et al. (2006)
Cariaco Basin	4.4–5.5	11.7 (200–225 m)	N.G.	Hashimoto et al. (1983)
Saanich Inlet	11.3–11.6	20.4 (110 m)	N.G.	Cohen (1978)

eastern South Pacific along latitude 32.5° S (Charpentier et al., 2010) also shows a very large increase (by a factor of 30) in N<sub>2</sub>O emissions within the coastal zone off Chile.

#### 4.5 European coastal systems

Seasonally-occurring major hypoxic/anoxic events have been reported from the Adriatic Sea (see e.g. Druon et al., 2004), the coastal Baltic Sea (see e.g. Conley et al., 2007) and the northwestern shelf of the Black Sea (see e.g. Daskalov, 2003). Persistent anoxia exists in the subsurface and deep waters of the Black Sea (see e.g. Kononov et al., 2005) and the basins of the central Baltic Sea (see e.g. Conley et al., 2009; BACC Author Team, 2008), discussed in detail in Sect. 5. A comprehensive overview of CH<sub>4</sub> and N<sub>2</sub>O measurements in European coastal waters is given in Bange (2006b). Within the context of this article we focus on the (rare) studies of CH<sub>4</sub> and N<sub>2</sub>O during hypoxic/anoxic events in shallow waters.

##### 4.5.1 Methane

Accumulation of CH<sub>4</sub> (up to 8 μM) in the Kiel Harbour has been observed during stagnation periods when the water column turns anoxic toward the end of the summer (Schmaljohann, 1996). Since June 2006, CH<sub>4</sub> measurements have been performed at the Boknis Eck Time Series Station (Eckernförde Bay, Southwestern Baltic Sea) on a monthly basis (Bange et al., 2010). Methane concentrations in the bottom layer (20–25 m) at Boknis Eck were up to 235 nM in October 2007; however, maximum CH<sub>4</sub> concentrations were not concurrent with hypoxic/anoxic events which usually occur in September/October. This apparent decoupling of hypoxic/anoxic events and the variability of CH<sub>4</sub> concentrations in the water column was explained by CH<sub>4</sub> release from sediments which occurred with a time lag of about one month after the sedimentation of organic material originating from phytoplankton blooms. Hypoxic/anoxic events seemed to have only a modulating effect on the accumulation of CH<sub>4</sub> in the water column (Bange et al., 2010). Considerable CH<sub>4</sub> release from sediments takes place despite the fact that CH<sub>4</sub> diffusion into the water column can be efficiently prevented by anaerobic CH<sub>4</sub> oxidation in the sediments of Eckernförde Bay (Treude et al., 2005).

##### 4.5.2 Nitrous oxide

At the Boknis Eck Time Series Station (Eckernförde Bay, SW Baltic Sea), N<sub>2</sub>O has been measured on a monthly basis since July 2005 (Schweiger, 2006; Bergmann, 2009). Figure 8 shows the temporal N<sub>2</sub>O variability during the hypoxic/anoxic event from September to October 2005. It is obvious that a pronounced N<sub>2</sub>O accumulation occurred after the water column was ventilated in November 2005. An increase of the N<sub>2</sub>O concentrations from 1–10 nM (in October 2005) to 16–20 nM (in November 2005) was recorded (Fig. 8).

Concurrent measurements of hydroxylamine (NH<sub>2</sub>OH), the first intermediate of NH<sub>4</sub><sup>+</sup> oxidation during nitrification, showed that maximum concentrations of NH<sub>2</sub>OH were observable in November 2005 as well. This led to the conclusion that both N<sub>2</sub>O and NH<sub>2</sub>OH were formed during the re-establishment of nitrification after the re-oxygenation of the anoxic waters at Boknis Eck (Schweiger et al., 2007).


#### 4.6 Chinese coastal waters

Because of recent economic development and population increase in Asia, coastal ecosystems in the region are subject to severe stress. Large rivers of the East and South Asian Rim are characterized by high nutrient concentrations with a skewed N/P ratio, owing most likely to extensive application of chemical fertilizers over the watersheds to support food production. As a result, depletion of dissolved O<sub>2</sub> commonly occurs in near-bottom coastal waters of Southeast Asia including the delta of the Mekong River (63–94 μM O<sub>2</sub>) and the Pearl River Estuary (31–63 μM O<sub>2</sub>) (Yin et al., 2004). Similar conditions have also been reported recently from the East China Sea (ECS) off the Changjiang Estuary (e.g. Fig. 9e), with bottom-water O<sub>2</sub> concentrations of 63–94 μM occurring over an area of 15 000 km<sup>2</sup> in August 1999 (Li et al., 2002).

The Changjiang empties into the ECS with a mean water discharge of  $928 \times 10^9 \text{ m}^3 \text{ a}^{-1}$  and total suspended matter (TSM) load of  $0.35 \times 10^9 \text{ t a}^{-1}$ , of which 70–80% occurs during the flood season (May–October) (Yang et al., 2002). The freshwater plume from the Changjiang disperses at the surface in the ECS (Fig. 9d) and at its peak in summer covers an area of ~85 000 km<sup>2</sup>. In winter, the Changjiang plume is restricted to the western side of the ECS, moving southward along the Chinese coast to the South China Sea. Extensive water exchange between ECS and Kuroshio occurs across the shelf break through upwelling and frontal processes. In the broad shelf region, patchy distribution of hydrographic properties is related to distinctive source water masses (cf. Su, 1998).

##### 4.6.1 Methane

Methane distribution in the ECS (e.g. Fig. 9a) shows a concentration gradient from the Changjiang Estuary to the Kuroshio Surface Waters with considerable seasonal and inter-annual variations (Zhang et al., 2008a). Elevated CH<sub>4</sub> concentrations occur in the water column of inner and mid-shelf but outside the high-turbidity plumes from the Changjiang, corresponding to the phytoplankton blooms caused by a decrease in turbidity that seems to limit photosynthesis closer to the coast (Zhang et al., 2004). Further offshore over the mid-shelf in ECS, where the water column is more stratified, CH<sub>4</sub> content of near-bottom samples can be 50–100% higher than at the surface. In the deeper water, data from interior of the Kuroshio show CH<sub>4</sub> concentrations



**Fig. 8.** Variability of N<sub>2</sub>O (A) and O<sub>2</sub> (B) at Boknis Eck time series station in the western Baltic Sea from July 2005 to May 2006. The hypoxic/anoxic event is marked by vertical dashed lines. Concentrations at 1 m and 25 m are highlighted with bold black and red lines respectively. The thin dashed lines represent sampling depths of 5, 10, 15 and 20 m. The arrow marks the measurements made after the ventilation event in November 2005. Data are from Schweiger (2006).

declining with depth down to 1000 m (Zhang et al., 2004). Over the ECS shelf, the upwelled Kuroshio subsurface waters have relatively low O<sub>2</sub> (125–156 μM) with lower CH<sub>4</sub> concentrations than in near-bottom waters over the inner shelf. Thus, in the coastal waters of the ECS that are affected by eutrophication, an increase of CH<sub>4</sub> saturation occurs in the water column (Fig. 9a), thereby sustaining a higher sea-to-air flux of CH<sub>4</sub> (2.5–5.0 μmol m<sup>-2</sup> d<sup>-1</sup>) relative to oligotrophic waters over the open-shelf (0.5–2.0 μmol m<sup>-2</sup> d<sup>-1</sup>) (Zhang et al., 2004). As stated above, hypoxia develops off the Changjiang Estuary (Fig. 9e) every summer (from June to September), with the lowest O<sub>2</sub> levels recorded in near-bottom waters being 16–31 μM between 20–50 m isobaths in the late summer of 2003. Also, waters with O<sub>2</sub> concentrations ranging from 125 to 156 μM can be found at the surface over the inner-shelf of the ECS, resulting from the dispersal of plumes from the Changjiang and upwelling of offshore waters because of the combination of buoyancy and topographic effects. Concentration of CH<sub>4</sub> in near-bottom


hypoxic waters can be as high as 30–40 nM, in comparison with the typical range of values between 10 and 20 nM in late autumn and early spring when the near-shore water column tends to be well mixed (Zhang et al., 2008a).

Freshwater runoff by the Changjiang appears to make a significant contribution to the CH<sub>4</sub> budget of the region. The concentrations in the affected area range between 15 and 130 nM with an average of 71.6 nM, and there is a positive correlation between river runoff and CH<sub>4</sub> concentration (Zhang et al., 2008a). This is similar to the observations off the river mouths elsewhere in the ocean, described earlier. Heterotrophic respiration in sediments overlain by hypoxic waters has been estimated to consume O<sub>2</sub> at a rate of 30–70 mmol m<sup>-2</sup> d<sup>-1</sup>, while CH<sub>4</sub> efflux from the sediments has been quantified as 1.7–2.2 μmol m<sup>-2</sup> d<sup>-1</sup>; CH<sub>4</sub> and O<sub>2</sub> appear to be negatively correlated in near-bottom waters affected by hypoxia (Zhang et al., 2008a). High surface concentrations in this region support a high sea-to-air flux, reaching up to 250 μmol m<sup>-2</sup> d<sup>-1</sup>, which is 20 times higher than the efflux from the shelf mixed water (10–15 μmol m<sup>-2</sup> d<sup>-1</sup>) and 2–3 orders of magnitude higher than the efflux from the Kuroshio Current region (<0.5 μmol m<sup>-2</sup> d<sup>-1</sup> – Zhang et al., 2008b).

#### 4.6.2 Nitrous oxide

Distribution (Fig. 9b) and air-sea exchange of N<sub>2</sub>O in the coastal region off the Changjiang Estuary and further offshore in the ECS are regulated by the prevailing hydrographic conditions (Fig. 9c–e) and nitrogen cycling. For instance, the upwelled water of the Kuroshio at the shelf break is characterized by relatively low O<sub>2</sub> with slightly higher N<sub>2</sub>O concentrations (20–30 nM) than surface waters of the open shelf (10–20 nM) and lower values than in the region affected by river runoff (30–40 nM) (Fig. 9b; Zhang et al., 2008b). Close to the coast, hypoxic subsurface waters having high salinity and low temperature are capped by warmer, brackish water plumes from the Changjiang during the summer season (Fig. 9c, d). Concentrations of N<sub>2</sub>O in the hypoxic layer (O<sub>2</sub> < 63 μM – Fig. 9e) can be as high as 40–60 nM as compared to 10–20 nM in the overlying surface waters and 5–10 nM at the surface further offshore over the open shelf (Zhang et al., 2008b). Where coastal hypoxia does not occur, vertical profiles of N<sub>2</sub>O concentrations in the near-shore and open shelf regions are similar, hovering around 10 nM. Off the Changjiang Estuary, ΔN<sub>2</sub>O is positively correlated with AOU but in a non-linear way; there also exist positive correlations between N<sub>2</sub>O on the one hand and NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> on the other suggesting that N<sub>2</sub>O is mainly produced through nitrification (Zhang et al., 2008b).

The sea-to-air flux of N<sub>2</sub>O in the region of coastal hypoxia off the Changjiang Estuary is up to 50–100 μmol m<sup>-2</sup> d<sup>-1</sup>, an order of magnitude higher than the values for the shelf mixed water of the ECS (<10 μmol m<sup>-2</sup> d<sup>-1</sup>), Kuroshio waters (<5 μmol m<sup>-2</sup> d<sup>-1</sup>) (Zhang et al., 2008b), and also the



**Fig. 9.** Distribution of hydrographic and chemical properties in the East China Sea from a station outside Changjiang Estuary (Y5) to the Korea Strait (Y1) in September 2003: (a) CH<sub>4</sub> (nM), (b) N<sub>2</sub>O (nM), (c) temperature (°C), (d) salinity, and (e) dissolved O<sub>2</sub> (μM). Modified from Zhang et al. (2008b).

Changjiang water ( $20 \mu\text{mol m}^{-2} \text{d}^{-1}$ , Jing Zhang, unpublished data). Even higher fluxes to the atmosphere are expected to occur during periods when stratification is broken down by vertical mixing. Moreover, high CH<sub>4</sub> and N<sub>2</sub>O concentrations can also be exported to the open ocean by circulation and dynamic processes across the shelf break, an important component of the “continental shelf pump” (Tsunogai et al., 1999).


#### 4.7 Gulf of Mexico

The biogeochemistry of the northern Gulf of Mexico is greatly affected by runoff from the Mississippi River, which is the sixth largest river in the world in terms of freshwater discharge ( $580 \text{ km}^3 \text{ a}^{-1}$  – Milliman and Meade, 1983). The enormous loading of nutrients and organic matter from land by the river in conjunction with strong near-surface stratification results in seasonal formation of a hypoxic zone that

seems to be expanding with time, presently occupying an area of  $22\,000 \text{ km}^2$  at its peak during summer (Rabalais and Turner, 2006). This region is also distinguished by the occurrence of hydrocarbon seeps and gas hydrates at the seafloor, additional factors that make it important for CH<sub>4</sub> cycling (Brooks et al., 1981).

##### 4.7.1 Methane

Supersaturation of surface waters of the Gulf of Mexico with respect to atmospheric CH<sub>4</sub> has long been known (Swinnerton and Lamontagne, 1974; Brooks et al., 1981). The occurrence of several subsurface maxima in the vertical profiles points to their multiple sources/formative processes such as release of CH<sub>4</sub> (originating from anoxic degradation of organic matter and from seeps or gas hydrates) from shelf sediments, lateral dispersal of CH<sub>4</sub>-rich layers, and in situ production (Brooks et al., 1981). An extensive data set on



**Fig. 10.** Vertical profiles of CH<sub>4</sub> at three stations located over the shelf in the region affected by the Mississippi plume in the Gulf of Mexico. Modified from Kelley (2003).

CH<sub>4</sub> in the Gulf of Mexico hypoxic zone has been generated by Kelley (2003) on a series of cruises conducted between March 1994 and July 1998. Her results show very high CH<sub>4</sub> concentrations, some even matching those observed off Namibia. These high values persist during all seasons, but tend to be more elevated during summer. Although surface waters had lower salinity (down to 20) as compared to bottom waters (~35), peak values were not always associated with the Mississippi plume with CH<sub>4</sub> maxima occurring at the surface (Fig. 10a), at mid-depth (within the pycnocline,

Fig. 10b) or close to the bottom (Fig. 10c). Kelley did not present O<sub>2</sub> data, if they were collected, but bottom-water hypoxia might have been present during the summer samplings (e.g. in case of Fig. 10b and c). The observed accumulation of CH<sub>4</sub> was ascribed to in situ production as a result of high concentration of particulate matter. Incubation of samples in the presence of picolinic acid (which inhibits CH<sub>4</sub> oxidation) provided evidence for in situ production. However, as in case of the Indian shelf, it is quite likely that the very high values (particularly at the surface) originate from river runoff. Moreover, emission of CH<sub>4</sub> from the sediments coupled with low O<sub>2</sub> content of the bottom water may also be important contributors to the greatly elevated CH<sub>4</sub> levels in the water column. In addition to the role of in situ production favoured by Kelley, mid-depth maxima can also be maintained by lateral mixing/advection of high-CH<sub>4</sub> water and/or the loss of CH<sub>4</sub> from the surface layer through air-sea exchange.

#### 4.7.2 Nitrous oxide

The first data set on N<sub>2</sub>O distribution and atmospheric emission from the Gulf of Mexico has just been made available by Walker et al. (2010). Their observations in August 2008 were interrupted by Tropical Storm Edouard, allowing them to quantify the emission pulse caused by storm-driven vertical mixing. Although hypoxic conditions prevailed in near bottom waters prior to the storm, the O<sub>2</sub> concentrations were not low enough (suboxic) for denitrification to occur. N<sub>2</sub>O concentrations ranged from 5 to 30 nM with the highest values observed at mid-depths or close to the bottom. These concentrations and the computed sea-to-air fluxes (0.6–6.9  $\mu\text{mol m}^{-2} \text{d}^{-1}$ ) were higher than those in the Caribbean and western Tropical Atlantic. The storm led to generally higher values of N<sub>2</sub>O (7–47 nM) as well as an increase in the N<sub>2</sub>O emission rate (1.2–11.7  $\mu\text{mol m}^{-2} \text{d}^{-1}$ ). While the higher surface concentrations were obviously due to entrainment of N<sub>2</sub>O from the hypoxic bottom water, the enhanced production through nitrification was attributed to reoxygenation of the water column and redistribution of organic nitrogen.

As in other areas (e.g. Rhee et al., 2009), inputs of N<sub>2</sub>O by river water to the Gulf of Mexico do not seem to be important as evident from the data of Fox et al. (1987) from the Mississippi Estuary. These data, collected during a period of low flow (October 1983), show close-to-saturation concentrations above a salinity of 20, and an almost conservative mixing behaviour in fresher waters with ~200% saturation occurring at 0 salinity.

#### 4.8 Tokyo Bay

A study of N<sub>2</sub>O was carried out by Hashimoto et al. (1999) in the Tokyo Bay where hypoxia is known to develop during summer. Observations on five cruises undertaken during



May–October, 1994, at a number of stations yielded high surface concentrations (averaging 11.8–90.3 nM) and saturations (166–1190%). However, these high values were apparently not related to O<sub>2</sub> depletion that prevailed in bottom waters during summer; instead their origin was a sewage treatment plant that generated a large amount of N<sub>2</sub>O (~0.1 Gg N a<sup>-1</sup>). In fact, the N<sub>2</sub>O concentration decreased with depth as the bottom waters were reducing and had lost almost all NO<sub>3</sub><sup>-</sup>. Anomalously high NO<sub>2</sub><sup>-</sup> or N<sub>2</sub>O values were thus not recorded, but since the gradients were sharp, it is possible that these features could have been missed due to inadequate sampling spacing in the vertical and with time. Once oxic conditions were restored throughout the water column in October, both NO<sub>3</sub><sup>-</sup> and N<sub>2</sub>O increased in bottom water but still remained well below the surface values.

#### 4.9 Chesapeake Bay


The Chesapeake Bay, the largest estuary in the United States and one of the largest in the world, has been experiencing seasonal hypoxia for several decades (e.g. Hagy et al., 2004; Kemp et al., 2009). Elkins et al. (1978) investigated vertical N<sub>2</sub>O distribution in the deeper parts of the estuary during a period of peak hypoxia (in July 1977). As in case of the Tokyo Bay, the suboxic sub-thermocline waters having very low NO<sub>3</sub><sup>-</sup> and NO<sub>2</sub><sup>-</sup> but high NH<sub>4</sub><sup>+</sup> concentrations were also depleted with N<sub>2</sub>O. Surface waters were moderately supersaturated during stable conditions but became slightly undersaturated following a mixing event.

### 5 Methane and nitrous oxide in enclosed anoxic basins

Land-locked or semi-enclosed basins with shallow sills connecting them with the open ocean (either directly or via another semi-enclosed sea), and having a circulation where deep water flows into the basin over the sill and the surface water flows out, experience either permanent (Black Sea and Cariaco Basin) or intermittent (Baltic Sea and Saanich Inlet) anoxia. All four examples listed here have distinct albeit narrow (several meters thick) suboxic layers that separate deep sulphidic waters (found below the sill depth) from the oxic layers lying above it. Steep chemical gradients (chemoclines) are conspicuous features of the suboxic zone (Cohen, 1978; Walter et al., 2006; Westley et al., 2006).

#### 5.1 Methane

One striking aspect of the biogeochemistry of these basins is the large accumulation of CH<sub>4</sub> in sulphidic waters (e.g. up to ~13 μM in the Black Sea (Fig. 11), 17 μM in the Cariaco Basin (Fig. 11) and 1.6 μM in the Saanich Inlet) (Bullister et al., 1982; Kessler et al., 2006a, b; Reeburgh, 2007; Ward et al., 1989). The source of CH<sub>4</sub> in these basins is believed to be sedimentary (Reeburgh, 2007). Kessler et al. (2005) measured <sup>14</sup>C in CH<sub>4</sub> in the Cariaco Basin and reported that




**Fig. 11.** Typical vertical profiles of CH<sub>4</sub> in the Black Sea (black symbols and curve) and the Cariaco Basin (red symbols and curve). Data are from Kessler et al. (2005, 2006a, b).

essentially all of the CH<sub>4</sub> in deep waters must be from fossil sources, in spite of earlier estimates of biogenic production (Reeburgh, 1976). However, deep-water CH<sub>4</sub> concentrations change relatively slowly with time in the Black Sea and Cariaco Basin, while data from the Saanich Inlet show more variability related to renewal of deep water (Ward et al., 1989). Given the high deep-water CH<sub>4</sub> levels, surface concentrations in these basins are elevated, but not abnormally so. Cariaco Basin surface water is at or slightly above saturation with the atmosphere (Ward et al., 1987; M. I. Scranton, unpublished), while in the Black Sea, where there are several seeps, high surface CH<sub>4</sub> can be observed (e.g. 294% saturation observed over a 90 m deep seep – Schmale et al., 2005).

#### 5.2 Nitrous oxide

Vertical distribution of N<sub>2</sub>O in anoxic basins is significantly different from that observed in the open-ocean suboxic zones. As expected, the sulphidic layer is devoid of N<sub>2</sub>O, a feature first noticed by Cohen (1978) in the Saanich Inlet (Fig. 12), and observed subsequently in other basins as well (e.g. Hashimoto et al., 1983; Butler and Elkins, 1991; Walter et al., 2006; Westley et al., 2006). Cohen (1978) found N<sub>2</sub>O concentration to rise up to 20.4 nM at the lower oxycline, followed by a rapid decline within the suboxic zone (<15 m thick) to reach zero/near-zero values at or just below the suboxic/anoxic interface (Fig. 12). The highest concentration



**Fig. 12.** Vertical profiles of (a) density ( $\sigma_t$ ) and N<sub>2</sub>O; (b) O<sub>2</sub> and H<sub>2</sub>S; (c) NH<sub>4</sub><sup>+</sup> and NO<sub>2</sub><sup>-</sup>; and (d) PO<sub>4</sub><sup>3-</sup> and NO<sub>3</sub><sup>-</sup> at two stations in the Saanich Inlet. Reproduced with permission from Cohen (1978).

is much lower than observed in open-ocean O<sub>2</sub>-deficient systems. But subsequent work has shown that this is not unusual for anoxic basins. A similar pattern of variability has been reported from the Cariaco Basin by Hashimoto et al. (1983), and from the Black Sea by Butler and Elkins (1991), and Westley et al. (2006). In fact, the peak N<sub>2</sub>O concentrations in these basins are even lower (<12 nM and <15 nM, respectively).

The spatial and temporal patterns of N<sub>2</sub>O variability observed in the Baltic Sea are quite puzzling. The first N<sub>2</sub>O data from this region were obtained when oxic conditions prevailed in the deep basins of the central Baltic Sea after a strong North Sea water inflow event in August/September 1977 (Rönnner, 1983). In July 1979, Rönnner fortuitously observed a dramatic change in N<sub>2</sub>O concentrations at ~100 m depth at one station in the western Gotland Basin exactly when the bottom water became anoxic. This shift was associated with a decrease in N<sub>2</sub>O concentration from 1523 nM to 0 nM within 24 h.

In a more recent study, Walter et al. (2006) investigated water-column distributions of N<sub>2</sub>O at 26 stations in the southern and central Baltic Sea in October 2003 after another major North Sea water inflow event in January 2003. By the time the observations were made, the O<sub>2</sub> rich North Sea water had already ventilated the formerly anoxic deep waters of the eastern Gotland Basin but had not ventilated the deep waters of the western Gotland Basin. In the anoxic water masses of the western basins N<sub>2</sub>O concentrations were <2 nM. The peak N<sub>2</sub>O concentration in the overlying waters here was around 20 nM, similar to the Black Sea and Saanich Inlet. This is substantially lower than the maximum concentration of 31 nM recorded in the Bornholm Basin that had been ventilated with O<sub>2</sub>-rich North Sea water. Walter et al. (2006) concluded that the shift from anoxic to oxic conditions after the inflow event in January 2003 had led to significant N<sub>2</sub>O accumulation in the water column. However, the accumulated N<sub>2</sub>O was not immediately released to the atmosphere because of the presence of a permanent halocline.

Westley et al. (2007) provided data on isotopic composition of N<sub>2</sub>O from the Black Sea that are very intriguing in that the  $\delta^{15}\text{N}$  and  $\delta^{18}\text{O}$  exhibited opposite trends, contrary to observations in other O<sub>2</sub>-deficient systems (e.g. Kim and Craig, 1990; Yoshinari et al., 1997; Naqvi et al., 1998). Within the N<sub>2</sub>O-maximum layer at the lower oxycline, there was a slight depletion of  $^{15}\text{N}$  and slight enrichment of  $^{18}\text{O}$  relative to the upper-layer mean values of 7.6‰ vs. air and 44.2‰ vs. SMOW, respectively. Much more pronounced isotopic shifts occurred within the suboxic layer where  $\delta^{15}\text{N}$  fell to a minimum of -10.8‰ (the lowest ever reported from seawater) whereas  $\delta^{18}\text{O}$  rose to a maximum of 67.0‰. The isotopomeric data revealed a  $^{15}\text{N}$  site preference maximum that coincided with the  $\delta^{18}\text{O}$  maximum. These results were interpreted to infer that both production of N<sub>2</sub>O through oxidation of NH<sub>4</sub><sup>+</sup> diffusing upward from the anoxic layer and consumption of N<sub>2</sub>O through denitrification were occurring within the suboxic layer, producing both depletion and enrichment of the heavier isotopes, respectively. In case of  $^{15}\text{N}$ , depletion due to production probably overwhelmed enrichment due to consumption. However, Westley et al. did not rule out other processes such as anammox influencing N<sub>2</sub>O cycling in the Black Sea. The above results underscore our incomplete understanding of the mechanisms of N<sub>2</sub>O production and consumption in systems approaching and reaching complete anoxia.

## 6 Discussion

It is believed that most of the CH<sub>4</sub> accumulating in bottom waters is derived from anoxic sediments through diffusion and bubble ebullition. The O<sub>2</sub> content/redox state of bottom waters is expected to affect sedimentary CH<sub>4</sub> production, its escape to the overlying water column, and its oxidative loss in the upper sedimentary column and in water itself. Therefore, one may expect an enhancement of CH<sub>4</sub> concentrations in O<sub>2</sub>-depleted bottom waters. However, O<sub>2</sub>-depletion does not seem to be a necessary and sufficient condition for dissolved CH<sub>4</sub> accumulation in the water column. High CH<sub>4</sub> concentrations (hundreds of nanomolar) are sometimes found in well-oxygenated waters overlying organic-rich sediments, as in the well known case of the Cape Lookout Bight (Martens and Klump, 1980) where hypoxic conditions do not develop at any time of the year (Kelley, 2003). Conversely, the world's largest naturally-formed coastal hypoxic zone over the Indian shelf does not exhibit large CH<sub>4</sub> build-up. Nevertheless, as discussed above and summarized in Table 2a, most O<sub>2</sub>-deficient systems from which CH<sub>4</sub> data are available accumulate CH<sub>4</sub> to varying degrees. This includes the three major open-ocean OMZs in the ETNP (Fig. 5c), ETSP and the Arabian Sea where significantly elevated concentrations of CH<sub>4</sub> are found within the suboxic layer (Burke et al., 1983; Sansone et al., 2004; Kelley and Jeffrey, 2002; Jayakumar et al., 2001). The CH<sub>4</sub> maximum

coincides with the intermediate nepheloid layer, a ubiquitous feature of suboxic zones (Burke et al., 1983; Jayakumar et al., 2001), and so in addition to the continental margin origin (Sansone et al., 2001), in situ production within reducing microenvironments is another plausible mechanism of its formation.

Maximal CH<sub>4</sub> build-up (to micromolar levels) in seawater occurs in truly anoxic (sulphidic) enclosed basins. In the most prominent of such basins, having the highest CH<sub>4</sub> concentrations (the Black Sea and the Cariaco Basin), <sup>14</sup>CH<sub>4</sub> measurements have established that fossil CH<sub>4</sub> released at the seafloor dominates over that produced through sediment diagenesis as a source to the water column (Kessler et al., 2005, 2006a). Thus, while anoxic conditions probably facilitate CH<sub>4</sub> preservation, their contribution to its production may be less important. Methane concentrations also increase, to a smaller extent (tens of nanomolar levels), in some silled basins overlain by hypoxic waters (e.g. the Gulf of California basins), but not in others (e.g. the California Borderland basins). It may be noted that occasional CH<sub>4</sub> maxima observed in the Santa Barbara Basin and Santa Monica Basin, reaching up to 50 nM (Kessler et al., 2008), most probably originate from lateral supply of CH<sub>4</sub> seeping from the seafloor. The above differences among hypoxic silled basins may be attributed to differences in sedimentary CH<sub>4</sub> supply and the residence time of water below the sill.

Along the open coasts too, bottom-water CH<sub>4</sub> content responds variously to changes in the ambient O<sub>2</sub> concentration. Over the Namibian shelf, where CH<sub>4</sub> content is inversely related to O<sub>2</sub> concentration in the bottom water (Monteiro et al., 2006), anoxic (sulphidic) waters have the highest CH<sub>4</sub> concentrations reaching up to ~5 µM. The increase in concentration (to ~40–50 nM) over the western Indian shelf during periods of anoxia is two orders of magnitude smaller. As pointed out earlier, the switch-over from hypoxic to suboxic conditions has little effect on bottom-water CH<sub>4</sub> content in this region, possibly because of deeper sulphate-CH<sub>4</sub> transition in the sediment arising mainly from its low organic carbon content. In the hypoxic zone of the Gulf of Mexico, on the other hand, much higher (hundreds of nanomolar) CH<sub>4</sub> levels are maintained despite a less intense hypoxia, evidently due to much larger supply from the sediments. According to Bange et al. (2010), CH<sub>4</sub> production in sediments of the Eckernförde Bay (Baltic Sea) and its supply to bottom waters is primarily linked to phytoplankton blooms with some (~1 month) time lag, and hypoxia exerts only a secondary modulating effect. This may also apply to most other areas. This implies that in areas where CH<sub>4</sub> accumulation and bottom water O<sub>2</sub> deficiency appear to be closely linked (see Figs. 2 and 7), the two may be driven by the same phenomenon – local high primary production – rather than having a direct cause-effect relationship. The other extreme case, as exemplified by the Indian shelf, appears to be where the O<sub>2</sub> deficiency is primarily due to low O<sub>2</sub> content of source waters and the local primary production is not that high such

that bottom-water CH<sub>4</sub> and O<sub>2</sub> are, to a large extent, decoupled.

In hypoxic coastal areas located close to river mouths (e.g. Gulf of Mexico, East China Sea, eastern Arabian Sea and northern Bay of Bengal), CH<sub>4</sub> concentrations are also enhanced by riverine inputs. Another complexity arises from in situ production of CH<sub>4</sub> in the water column itself, for which there is considerable circumstantial evidence (Scranton and Brewer, 1977; Scranton and Farrington, 1977; Karl and Tilbrook, 1994; Kelley, 2003; Karl et al., 2008). This is reflected by significant supersaturation in surface water in productive margins that experience upwelling but no hypoxia (e.g. Kock et al., 2008). More often than not, however, hypoxia – both natural and human induced – occurs in regions of high productivity where in situ production rate in oxygenated water column is also expected to be high, and it is difficult to differentiate such production from enhanced emissions from the sediments (Scranton and Farrington, 1977).

Estimates of sea-to-air flux densities from major hypoxic/suboxic/anoxic zones are summarized in Table 2a for CH<sub>4</sub> and Table 2b for N<sub>2</sub>O. An effort has been made in Tables 3a and b, respectively for the two gases, to quantify total emissions from all areas of the oceans affected by O<sub>2</sub> deficiency using representative figures for various types of systems. However, while considering these estimates the following caveats must be kept in mind:

1. The estimated fluxes given in Table 2 were obtained using different models of the air-sea gas exchange.
2. Only diffusive fluxes are considered, ignoring emissions via bubble ebullition. The latter are significant in the case of CH<sub>4</sub>, even exceeding diffusive fluxes in some shallow regions. For instance, Martens and Klump (1980) measured a large flux of gaseous CH<sub>4</sub> (averaging 16.8 mmol m<sup>-2</sup> (low tide)<sup>-1</sup>) in summer out of the sediments of the Cape Lookout Bight. Approximately 85% of the rising bubbles escaped dissolution in this shallow setting, resulting in an annually-averaged emission of ~12 mmol CH<sub>4</sub> m<sup>-2</sup> d<sup>-1</sup> to the atmosphere. This flux is 5–7 times the combined inputs of CH<sub>4</sub> to the water column through diffusion from sediments and dissolution of bubbles. Assuming that ~1% of the global continental shelf area shallower than 10 m (1.4 × 10<sup>6</sup> km<sup>2</sup>) has a bubble ebullition CH<sub>4</sub> flux density comparable to that in the Cape Lookout Bight, an upper limit on the total flux from such systems could be placed at ~1 Tg a<sup>-1</sup>. Not much is known about the bubble ebullition flux from coastal hypoxic zones except that such flux should be important over the Namibian shelf (Brüchert et al., 2006, 2008), and also expected to be so in the Gulf of Mexico, given the high CH<sub>4</sub> concentration in the water column (e.g. Fig. 10c).

**Table 3a.** Total CH<sub>4</sub> emissions from various types of marine O<sub>2</sub>-deficient systems.

Region	Area (km <sup>2</sup> )	Flux density (μmol m <sup>-2</sup> d <sup>-1</sup> )	Total flux (Tg CH <sub>4</sub> a <sup>-1</sup> )
Open-ocean hypoxic zones	29.3 × 10 <sup>6</sup> <sup>a</sup>	1.6 <sup>b</sup> –2.2 <sup>c</sup>	0.27–0.38
Naturally-formed continental-margin hypoxic zones	1.1 × 10 <sup>6</sup> <sup>d</sup>	2.8 <sup>e</sup> –11.2 <sup>f</sup>	0.02–0.07
Athropogenically-formed hypoxic zones			
Persistent	88 000 <sup>h</sup>	21–36 <sup>i</sup>	0.01–0.02
Periodic/Seasonal <sup>g</sup>	138 000 <sup>h</sup>	21–36 <sup>i</sup>	0.008–0.015
Enclosed anoxic basins	0.83 × 10 <sup>6</sup>	10–50 <sup>j</sup>	0.05–0.24
Total	31.5 × 10 <sup>6</sup>		0.36–0.72

<sup>a</sup> Paulmier and Ruiz-Pino (2009); <sup>b</sup> Open Arabian Sea – Naqvi et al. (2005); <sup>c</sup> Open ETNP – Sansone et al. (2001); <sup>d</sup> Helly and Levin (2004); <sup>e</sup> Mexican continental margin – Sansone et al. (2004); <sup>f</sup> Western Indian shelf – Jayakumar (1999); <sup>g</sup> Assumed for 6 months; <sup>h</sup> Diaz and Rosenberg (2008); <sup>i</sup> East China Sea – Zhang et al. (2008); <sup>j</sup> Chosen from various references in Table 2a.

**Table 3b.** Total N<sub>2</sub>O emissions from various types of marine O<sub>2</sub>-deficient systems.

Region	Area (km <sup>2</sup> )	Flux density (μmol m <sup>-2</sup> d <sup>-1</sup> )	Total flux (Tg N <sub>2</sub> O a <sup>-1</sup> )
Open-ocean hypoxic zones	29.3 × 10 <sup>6</sup>	2.7 <sup>a</sup> –4.5 <sup>b</sup>	1.27–2.12
Naturally-formed continental-margin hypoxic zones	1.1 × 10 <sup>6</sup>	10–50 <sup>c</sup>	0.18–0.88
Athropogenically-formed hypoxic zones			
Persistent	88 000	3.3 <sup>d</sup> –17.1 <sup>e</sup>	0.004–0.024
Periodic/Seasonal <sup>f</sup>	138 000	3.3 <sup>d</sup> –17.1 <sup>e</sup>	0.003–0.019
Enclosed anoxic basins	0.83 × 10 <sup>6</sup>	1.6–5.2 <sup>g</sup>	0.02–0.07
Total	31.5 × 10 <sup>6</sup>		1.48–3.11

<sup>a</sup> ETNP – Cohen and Gordon (1978); <sup>b</sup> Open Arabian Sea – Naqvi and Noronha (1991); <sup>c</sup> Chosen from various references in Table 2b; <sup>d</sup> Gulf of Mexico – Walker et al. (2010); <sup>e</sup> East China Sea – Zhang et al. (2008); <sup>f</sup> Assumed for 6 months; <sup>g</sup> Black Sea – Amouroux et al. (2002).

The extent of CH<sub>4</sub> supply to the atmosphere by bubbles rising from seeps, clathrates and mud volcanoes at deeper seafloor in areas such as the Black Sea and Gulf of Mexico has become a subject of considerable interest in recent years. McGinnis et al. (2006) quantified dissolution of CH<sub>4</sub> from such bubbles from a combination of modelling and acoustic observations and concluded that most of the CH<sub>4</sub> carried by bubbles venting from sites deeper than 100 m in the Black Sea does not reach the atmosphere. However, a more recent study by Solomon et al. (2009) involving direct submersible sampling of water around hydrocarbon plumes in the Gulf of Mexico has challenged this view. These authors showed that bubble dissolution could be inhibited by bubble size, upwelling flows and the presence of surfactants. This

together with slow oxidation seems to ensure significant transport of CH<sub>4</sub> from deeper seep sites to the surface layer and atmosphere. Thus, the issue is yet to be fully settled.

3. Ranges of various flux density estimates provided in Table 2 are quite large. As a consequence, selection of representative values for various types of systems (Table 3) leads to large uncertainty. In most cases, we have chosen reasonable reported maximal and minimal mean flux density values so as to keep the ranges as narrow as possible. Another uncertainty pertains to the estimates of areas covered by various types of systems, especially those formed as a result of human activities (eutrophication). These estimates, from Diaz and Rosenberg (2008), are available for only one-quarter of

all anthropogenically-formed hypoxic zones. The remaining three-quarters of such systems are smaller in size, and hence would not add very much to the total area. Moreover, this may be compensated to some extent by the fact that a few systems falling under the anthropogenic group are also included elsewhere (e.g. the Danish coastal waters are counted again under enclosed anoxic basins). For these reasons, estimates of fluxes provided in Table 3 should be treated as crude first approximations, enabling an evaluation of their relative importance.

Despite modest sea-to-air flux density (Table 2a) the open-ocean OMZs are the largest source of CH<sub>4</sub> diffusing to the atmosphere from the sea surface among all hypoxic regions listed in Table 3a. This term is also the best constrained as estimates of the flux density for all other groups are much more variable. Among the naturally-formed hypoxic zones of the continental margins, the Namibian shelf is arguably the most important CH<sub>4</sub> emitter. However, atmospheric fluxes of CH<sub>4</sub> from this region are very poorly quantified. To give an idea of the potential importance of this region, an area of  $\sim 100\,000\text{ km}^2$  of the suboxic zone (Kuypers et al., 2005) and an assumed average flux density of  $200\,\mu\text{mol m}^{-2}\text{ d}^{-1}$  (toward the lower range of estimate in Table 2a for this region) would yield a total annual flux of  $\sim 0.12\text{ Tg CH}_4$ . This very crude calculation shows that the total CH<sub>4</sub> efflux from natural hypoxic zones of the continental margins may be comparable to that from open-ocean OMZs. The enclosed anoxic basins occupy slightly smaller area but are distinguished by the highest CH<sub>4</sub> concentrations. However, as the CH<sub>4</sub>-rich anoxic, subsurface waters are well isolated from the atmosphere, the total emission from these systems is not very large. Finally, at present there is a dearth of data on CH<sub>4</sub> emissions from anthropogenically-formed coastal hypoxic zones. If the estimates from the region off the Changjiang mouth in the East China Sea are taken to be representative of this group, as has been done in Table 3a, emissions from this group will be negligible. However, it should be noted that, given the high surface concentrations reported by Kelley (2003) from the Gulf of Mexico hypoxic zone, CH<sub>4</sub> flux density for this region, and in some other similar systems, may be substantially higher than that reported for the East China Sea. All marine areas affected by hypoxia put together emit up to  $\sim 1\text{ Tg CH}_4$  annually. While constituting a significant fraction of global oceanic CH<sub>4</sub> emission (in fact, it is comparable to the total CH<sub>4</sub> efflux from global ocean,  $0.6\text{--}1.2\text{ Tg a}^{-1}$ , reported by Rhee et al., 2010), it still represents merely 0.2% of all inputs of CH<sub>4</sub> to the atmosphere (Reeburgh, 2007).

In view of the lack of evidence for a primary control of bottom-water hypoxia on sedimentary CH<sub>4</sub> production, as well as low estimated emissions of CH<sub>4</sub> from the anthropogenically-formed coastal hypoxic zones, it is highly unlikely that an intensification or expansion of such zones

will have a large impact on global oceanic emissions of CH<sub>4</sub> to the atmosphere. A more important factor favouring methanogenesis in coastal sediments could be the potential increase in productivity due to eutrophication or intensification of coastal upwelling due to global warming (Bakun, 1990). At present, it is difficult to quantify or predict such changes. Nevertheless, it would be reasonable to conclude that unless CH<sub>4</sub> emissions are enhanced by more than an order of magnitude, the present status, where emissions from the ocean as a whole form an insignificant term in the atmospheric CH<sub>4</sub> budget, is unlikely to change.

The available information on N<sub>2</sub>O distribution and computed fluxes across the air-sea interface is summarized in Table 2b. Nitrous oxide cycling differs from that of CH<sub>4</sub> in three respects: (1) N<sub>2</sub>O is mainly produced in the water column; (2) its production by all known mechanisms is enhanced at low O<sub>2</sub> concentrations; and (3) it has no sink in oxygenated waters. As stated earlier, N<sub>2</sub>O does get reduced to N<sub>2</sub> within the cores of suboxic zones, but the peripheries of these zones provide most suitable conditions for its production. Consequently, O<sub>2</sub>-deficient aquatic systems are generally net producers of N<sub>2</sub>O (e.g. Codispoti et al., 1992). However, the extent to which the loss of O<sub>2</sub> promotes N<sub>2</sub>O production varies from one system to another (Table 2b) for reasons that are still not clear. The response of N<sub>2</sub>O to deoxygenation is non-linear. The “normal” behaviour is that N<sub>2</sub>O values peak at tens of nanomolar levels just before the environment turns reducing (e.g. Fig. 5a). Such enhancement of the N<sub>2</sub>O yield, recorded in both (largely) natural (e.g. off Oman) and (largely) anthropogenic (e.g. off China) hypoxic/suboxic systems (Table 2a), is most likely due to nitrifier denitrification (oxidation of NH<sub>4</sub><sup>+</sup> to NO<sub>2</sub><sup>-</sup> followed by the reduction of NO<sub>2</sub><sup>-</sup> to gaseous nitrogen species by autotrophic nitrifiers) (Yamagishi et al., 2007). Deviations from this behaviour, where N<sub>2</sub>O concentrations climb up to hundreds of nanomolar levels (Table 2b), have been observed in three naturally-formed coastal suboxic environments (off India, Peru and Chile). We predict similar abnormally high N<sub>2</sub>O concentrations off Namibia as well. The most important common feature of these systems is the extension of OMZ to very shallow depths. The most probable way by which N<sub>2</sub>O accumulates in very high concentrations in such settings is a more rapid reduction of NO<sub>2</sub><sup>-</sup>/NO to N<sub>2</sub>O as compared to its own reduction to N<sub>2</sub> by denitrifiers. This requires an inhibition of the activity of N<sub>2</sub>O reductase. Although it has been speculated that frequent aeration due to turbulence may deactivate this enzyme from time to time in shallow, rapidly denitrifying waters (Naqvi et al., 2000), the cause of such abrupt change in N<sub>2</sub>O cycling is not well understood and difficult to predict. This is, for example, illustrated by data from the Boknis Eck time series station (see Fig. 8) where N<sub>2</sub>O only peaks when the system is shifting from suboxic/anoxic (reducing) to oxic conditions which is in contrast to the “normal” behaviour discussed above.

Behaviour of N<sub>2</sub>O in anoxic basins is quite puzzling. While its reduction to N<sub>2</sub> expectedly accounts for near-zero levels in sulphidic waters, the build-up below the oxygenated mixed layer is well below expectation (e.g. Fig. 12), except for the aforementioned event in the Baltic where N<sub>2</sub>O concentration shot up to a world-record high level of  $\sim 1.5 \mu\text{M}$  (Table 2). One explanation for this could be that vertical transition from oxic to suboxic and then anoxic conditions in these basins is rapid, but then such is also the case over the western Indian shelf where, as stated above, abnormally high N<sub>2</sub>O concentrations often occur in close proximity to sulphidic waters. It would seem that shallow but unstable suboxia maintains high levels of N<sub>2</sub>O over the Indian shelf (Codispoti, 2010), whereas the nitrogen cycle is more often in a steady state in enclosed anoxic basins. Another important factor that probably contributes to the lower N<sub>2</sub>O levels in the anoxic basins is the low concentrations of secondary NO<sub>2</sub><sup>-</sup> (e.g. Fig. 12).

Among the anthropogenically-formed coastal hypoxic zones, N<sub>2</sub>O data are available from the East China Sea, Gulf of Mexico, Chesapeake Bay and Tokyo Bay. These systems show different trends that can be related to different redox conditions prevailing at the time of observations. That is, bottom waters were hypoxic in the East China Sea and Gulf of Mexico, and nearly-anoxic (having very low NO<sub>3</sub><sup>-</sup>+NO<sub>2</sub><sup>-</sup> and high NH<sub>4</sub><sup>+</sup>) in the other two areas. Accordingly, there was moderate accumulation of N<sub>2</sub>O at the first two sites. The Chesapeake Bay and Tokyo Bay exhibited a trend similar to enclosed anoxic basins, except for high surface values in the Tokyo Bay that clearly originated from a sewage treatment plant, as discussed earlier. Overall, the data gathered so far from the anthropogenically-formed coastal hypoxic zones have not shown the kind of N<sub>2</sub>O build-up that distinguishes naturally-formed, upwelling-related coastal suboxic zones, but a number of other anthropogenically-formed systems still remain to be investigated.

Estimates of sea-to-air flux density generally reflect the trends of N<sub>2</sub>O variability in the water column (Table 2b). The highest flux densities are computed for the naturally-formed hypoxic zones of the continental margins followed by open-ocean OMZs, anthropogenically-formed coastal hypoxic zones and enclosed anoxic zones in decreasing order. As in case of CH<sub>4</sub>, total efflux of N<sub>2</sub>O is the largest from open-ocean OMZs. This estimate also happens to be the most robust. Estimates for the naturally-formed coastal hypoxic zones are much more uncertain; even for the data-rich hypoxic zone over the Indian shelf, they vary from 0.05 to 0.38 Tg N<sub>2</sub>O a<sup>-1</sup> (Naqvi et al., 2006b). The upper limit of emissions estimated by us for this group (0.88 Tg N<sub>2</sub>O a<sup>-1</sup>) compares well with the reported global emission of 0.86 Tg N<sub>2</sub>O-N a<sup>-1</sup> from the estuaries and continental shelves by Seitzinger and Kroeze (1998). Our computations show that the anthropogenically-formed coastal hypoxic zones and enclosed basins presently do not contribute very significantly

to the total marine N<sub>2</sub>O emission to the atmosphere. Our estimate for N<sub>2</sub>O emission from all hypoxic/suboxic/anoxic zones (1.48–3.11 Tg N<sub>2</sub>O a<sup>-1</sup>) represents 16–33% of the total oceanic N<sub>2</sub>O source to the atmosphere, lower than 50% considered by Codispoti (2010). However, it is higher than the estimate for efflux of N<sub>2</sub>O from the global oceans (0.9–1.7 Tg N<sub>2</sub>O-N a<sup>-1</sup>) provided by Rhee et al. (2009). Considering such a substantial contribution from areas experiencing O<sub>2</sub> deficiency in the water column to the global marine N<sub>2</sub>O emission, and of the latter to the atmospheric N<sub>2</sub>O budget, and also taking into account the sensitivity of N<sub>2</sub>O cycling in aquatic systems to minor changes in the ambient O<sub>2</sub> concentration in the very low range, one may conclude that any major alterations in O<sub>2</sub> distribution in coastal as well as off-shore waters are certain to have a large impact on the N<sub>2</sub>O budget. However, this effect cannot be quantified at present because of continuing uncertainties concerning the basic formative mechanisms as well as a lack of observations in key coastal regions.

**Acknowledgements.** This manuscript was prepared during the tenure of a Marie Curie Incoming International Fellowship awarded to SWAN, who expresses his gratitude to the European Commission and his host Marcel Kuypers, Director, Max-Planck Institute for Marine Microbiology, Bremen, for support and facilities. The authors thank Alina Freing for drafting Fig. 1. They are also grateful to Frank Sansone and John Kessler for making available data used in Figs. 6 and 7, and Fig. 11, respectively. Thanks are also due to Ed Urban and Denis Gilbert for their keen interest and constant encouragement. The paper has immensely benefitted by the reviews provided by Gwenaél Abril and another anonymous referee.

The service charges for this open access publication have been covered by the Max Planck Society.

Edited by: D. Gilbert

## References

- Anonymous: Greenhouse gases hit modern-day highs, *Nature*, 456, 558–559, 2008.
- Amouroux, D., Roberts, G., Rapsomanikis, S., and Andreae, M. O.: Biogenic gas (CH<sub>4</sub>, N<sub>2</sub>O, DMS) emission to the atmosphere from near-shore and shelf waters of the north-western Black Sea, *Estuar. Coast. Shelf Sci.*, 54, 575–587, 2002.
- BACC (BALTEX Assessment of Climate Change) Author Team: Assessment of Climate Change for the Baltic Sea Basin, Springer, Berlin, 474 pp., 2008.
- Bakun, A.: Global climate change and intensification of coastal ocean upwelling, *Science*, 247, 198–201, 1990.
- Bange, H. W.: New Directions: The importance of the oceanic nitrous oxide emissions, *Atmos. Environ.*, 40, 198–199, 2006a.
- Bange, H. W.: Nitrous oxide and methane in European coastal waters, *Estuar. Coast. Shelf Sci.*, 70, 361–374, 2006b.
- Bange, H. W.: Gaseous nitrogen compounds (NO, N<sub>2</sub>O, N<sub>2</sub>, NH<sub>3</sub>) in the ocean, in: *Nitrogen in the Marine Environment*, 2nd Edi-



- tion, edited by: Capone, D. G., Bronk, D. A., Mulholland, M. R., and Carpenter, E. J., Elsevier, Amsterdam, 51–94, 2008.
- Bange, H. W., Bartell, U. H., Rapsomanikis, S., and Andreae, M. O.: Methane in the Baltic and North Seas and a reassessment of the marine emissions of methane, *Global Biogeochem. Cycles*, 8, 465–480, 1994.
- Bange, H. W., Rapsomanikis, S., and Andreae, M. O.: Nitrous oxide emissions from the Arabian Sea, *Geophys. Res. Lett.*, 23, 3175–3178, 1996.
- Bange, H. W., Ramesh, R., Rapsomanikis, S., and Andreae, M. O.: Methane in surface waters of the Arabian Sea, *Geophys. Res. Lett.*, 25, 3547–3550, 1998.
- Bange, H. W., Andreae, M. O., Lal, S., Law, C. S., Naqvi, S. W. A., Patra, P. K., Rixen, T., and Upstill-Goddard, R. C.: Nitrous oxide emissions from the Arabian Sea: A synthesis, *Atmos. Chem. Phys.*, 1, 61–71, doi:10.5194/acp-1-61-2001, 2001a.
- Bange, H. W., Rapsomanikis, S., and Andreae, M. O.: Nitrous oxide cycling in the Arabian Sea, *J. Geophys. Res.*, 106, 1053–1066, 2001b.
- Bange, H. W., Bergmann, K., Hansen, H. P., Kock, A., Koppe, R., Malien, F., and Ostrau, C.: Dissolved methane during hypoxic events at the Boknis Eck time series station (Eckernförde Bay, SW Baltic Sea), *Biogeosciences*, 7, 1279–1284, doi:10.5194/bg-7-1279-2010, 2010.
- Bates, T. S., Kelly, K. C., Johnson, J. E., and Gammon, R. H.: A reevaluation of the open ocean source of methane to the atmosphere, *J. Geophys. Res.*, 101, 6953–6961, 1996.
- Bergmann, K.: Spurengasmessungen an der Zeitserienstation Boknis Eck, Diploma thesis, Kiel University, Kiel, 75 + XIII pp., 2009.
- Berner, U., Poggenburg, J., Faber, E., Quadfasel, D., and Frische, A.: Methane in ocean waters of the Bay of Bengal: its sources and exchange with the atmosphere, *Deep-Sea Res. II*, 50, 925–950, 2003.
- Biswas, H., Mukhopadhyay, S. K., Sen, S., and Jana, T. K.: Spatial and temporal patterns of methane dynamics in the tropical mangrove dominated estuary, NE coast of Bay of Bengal, India, *J. Mar. Sys.*, 68, 55–64, 2007.
- Bograd, S. J., Castro, C. G., Di Lorenzo, E., Palacios, D. M., Bailey, H., Gilly, W., and Chavez, F. P.: Oxygen declines and the shoaling of the hypoxic boundary in the California Current, *Geophys. Res. Lett.*, 35, L12607, doi:10.1029/2008GL034185, 2008.
- Brettar, I. and Rheinheimer, G.: Denitrification in the central Baltic: Evidence for H<sub>2</sub>S oxidation as motor of denitrification at the oxic-anoxic interface, *Mar. Ecol. Prog. Ser.*, 77, 157–169, 1991.
- Brooks, J. M., Reid, D. F., and Bernard, B. B.: Methane in the upper water column of the Northwestern Gulf of Mexico, *J. Geophys. Res.*, 86, 11029–11040, 1981.
- Brüchert, V., Currie, B., Peard, K. R., Lass, U., Endler, R., Dübecke, A., Julies, E., Leipe, T., and Zitzmann, S.: Biogeochemical and physical control on shelf anoxia and water column hydrogen sulphide in the Benguela upwelling system off Namibia, in: *Past and Present Water Column Anoxia*, edited by: Neretin, L. N., NATO Science Series, IV. Earth and Environmental Sciences – vol. 64, Springer, Dordrecht, 161–193, 2006.
- Brüchert, V., Currie, B., and Peard, K. R.: Hydrogen sulphide and methane emissions on the central Namibian shelf, *Prog. Oceanogr.*, 83, 169–179, 2009.
- Bullister, J. L., Guinasso Jr., N. L., and Schink, D. R.: Dissolved hydrogen, carbon monoxide, and methane at the CEPEX site, *J. Geophys. Res.*, 87, 2022–2034, 1982.
- Burke Jr., R. A., Reid, D. F., Brooks, J. M., and Lavoie, D. M.: Upper water column methane geochemistry in the eastern tropical North Pacific, *Limnol. Oceanogr.*, 28, 19–32, 1983.
- Bussmann, I. and Suess, E.: Groundwater seepage in Eckernförde Bay (Western Baltic Sea): Effect on methane and salinity distribution of the water column, *Cont. Shelf Res.*, 18, 1795–1806, 1998.
- Butler, J. H. and Elkins, J. W.: An automated technique for the measurement of dissolved N<sub>2</sub>O in natural waters, *Mar. Chem.*, 34, 47–61, 1991.
- Butler, J. H., Elkins, J. W., Thompson, T. M., and Egan, K. B.: Tropospheric and dissolved N<sub>2</sub>O of the West Pacific and East Indian Oceans during the El Niño Southern Oscillation event of 1987, *J. Geophys. Res.*, 94, 14865–14877, 1989.
- Carr, M.-E.: Estimation of potential productivity in eastern boundary currents using remote sensing, *Deep-Sea Res. II*, 49, 59–80, 2002.
- Chan, F., Barth, J. A., Lubchenco, J., Kirincich, A., Weeks, H., Peterson, W. T., and Menge, B. A.: Emergence of anoxia in the California current large marine ecosystem, *Science*, 319, 920–920, 2008.
- Chapman, P. and Shannon, L. V.: The Benguela Ecosystem. Part II: chemistry and related processes, *Oceanogr. Mar. Biol. Annu. Rev.*, 23, 183–251, 1985.
- Charpentier, J., Farias, L., and Pizarro, O.: Nitrous oxide fluxes in the Central and Eastern South Pacific, *Global Biogeochem. Cycles*, in press, 2010.
- Cline, J. D. and Richards, F. A.: Oxygen deficient conditions and nitrate reduction in the eastern tropical North Pacific Ocean, *Limnol. Oceanogr.*, 17, 885–900, 1972.
- Cline, J. D. and Kaplan, I. R.: Isotopic fractionation of dissolved nitrate during denitrification in the eastern tropical North Pacific Ocean, *Mar. Chem.*, 3, 271–299, 1975.
- Codispoti, L. A.: Interesting times for marine N<sub>2</sub>O, *Science*, 327, 1339–1340, 2010.
- Codispoti, L. A. and Christensen, J. P.: Nitrification, denitrification and nitrous oxide cycling in the eastern tropical South Pacific Ocean, *Mar. Chem.*, 16, 277–300, 1985.
- Codispoti, L. A., Friederich, G. E., Packard, T. T., Glover, H. E., Kelly, P. J., Spinrad, R. W., Barber, R. T., Elkins, J. W., Ward, B. B., Lipschultz, F., and Lostaunau, N.: High nitrite levels off Northern Peru: A signal of instability in the marine denitrification rate, *Science*, 233, 1200–1202, 1986.
- Codispoti, L. A., Elkins, J. W., Friederich, G. E., Packard, T. T., Sakamoto, C. M., and Yoshinari, T.: On the nitrous oxide flux from productive regions that contain low oxygen waters, in: *Oceanography of the Indian Ocean*, edited by: Desai, B. N., Oxford-IBH, New Delhi, 271–284, 1992.
- Codispoti, L. A., Brandes, J. A., Christensen, J. P., Devol, A. H., Naqvi, S. W. A., Paerl, H. W., and Yoshinari, T.: The oceanic fixed nitrogen and nitrous oxide budgets: Moving targets as we enter the Anthropocene?, *Sci. Mar.*, 65, 85–105, 2001.
- Cohen, Y.: Consumption of dissolved nitrous oxide in an anoxic basin, Saanich Inlet, British Columbia, *Nature*, 272, 235–237, 1978.
- Cohen, Y. and Gordon, L. I.: Nitrous oxide in oxygen minimum of eastern tropical North Pacific – Evidence for its consumption

- during denitrification and possible mechanisms for its production, *Deep-Sea Res.*, 25, 509–524, 1978.
- Conley, D. J., Carstensen, J., Aertebjerg, G., Christensen, P. B., Dalsgaard, T., Hansen, J. L. S., and Josefson, A. B.: Long-term changes and impacts of hypoxia in Danish coastal waters, *Ecol. Appl.*, 17, S165–S184, 2007.
- Conley, D. J., Björck, S., Bonsdorff, E., Carstensen, J., et al.: Hypoxia-related processes in the Baltic Sea, *Environ. Sci. Technol.*, 43, 3412–3420, 2009.
- Connolly, T. P., Hickey, B. M., Geier, S. L., and Cochlan, W. P.: Processes influencing seasonal hypoxia in the northern California Current System, *J. Geophys. Res.*, 115, C03021, doi:10.1029/2009JC005283, 2010.
- Copenhagen, W. J.: The periodic mortality of fish in the Walvis region – a phenomenon within the Benguela Current, *Investigational Report Division of Fisheries-Union of South Africa*, 14, 1–35, 1953.
- Cornejo, M., Farías, L., and Paulmier, A.: Temporal variability in N<sub>2</sub>O water content and air-sea exchange in an upwelling area off central Chile (36° S), *Mar. Chem.*, 101, 85–94, 2006.
- Cornejo, M., Farías, L., and Gallegos, M.: Seasonal cycle of N<sub>2</sub>O vertical distribution and air-sea fluxes over the continental shelf waters off central Chile (~36° S), *Prog. Oceanogr.*, 75, 383–395, 2007.
- Cynar, F. J. and Yayanos, A. A.: The distribution of methane in the upper waters of the Southern California Bight, *J. Geophys. Res.*, 97, 11269–11285, 1992.
- Cynar, F. J. and Yayanos, A. A.: The oceanic distribution of methane and its flux to the atmosphere over Southern California waters, in: *Biogeochemistry of Global Change – Radiatively Active Gases*, edited by: Oremland, R. S., Chapman and Hall, London, 551–573, 1993.
- Damm, E., Helmke, E., Thoms, S., Schauer, U., Nöthig, E., Bakker, K., and Kiene, R. P.: Methane production in aerobic oligotrophic surface water in the central Arctic Ocean, *Biogeosciences*, 7, 1099–1108, doi:10.5194/bg-7-1099-2010, 2010.
- Daneri, G., Dellarossa, V., Quiñones, R., Jacob, B., Montero, P., and Ulloa, O.: Primary production and community respiration in the Humboldt Current System off Chile and associated oceanic areas, *Mar. Ecol. Prog. Ser.*, 197, 41–49, 2000.
- Daskalov, G. M.: Long-term changes in fish abundance and environmental indices in the Black Sea, *Mar. Ecol. Prog. Ser.*, 255, 259–270, 2003.
- Deuser, W. G.: Reducing environments, in: *Chemical Oceanography*, vol. 3, edited by: Riley, J. P. and Chester, R., Academic Press, London, 1–37, 1975.
- de Wilde, H. P. J. and Helder, W.: Nitrous oxide in the Somali Basin: the role of upwelling, *Deep-Sea Res. II*, 44, 1319–1340, 1997.
- Diaz, R. J. and Rosenberg, R.: Spreading dead zones and consequences for marine ecosystems, *Science*, 321, 926–929, 2008.
- Dore, J. E., Popp, B. N., Karl, D. M., and Sansone, F. J.: A large source of atmospheric nitrous oxide from subtropical North Pacific surface waters, *Nature*, 396, 63–66, 1998.
- Druon, J.-N., Schrimpf, W., Dobricic, S., and Stips, A.: Comparative assessment of large-scale marine eutrophication: North Sea area and Adriatic Sea as case studies, *Mar. Ecol. Prog. Ser.*, 272, 1–23, 2004.
- Duce, R. A., LaRoche, J., Altieri, K., Arrigo, K. R., et al.: Impacts of atmospheric anthropogenic nitrogen on the open ocean, *Science*, 320, 893–897, 2008.
- Dugdale, R. C., Goering, J. J., Barber, R. T., Smith, R. L., and Packard, T. T.: Denitrification and hydrogen sulfide in Peru upwelling during 1976, *Deep-Sea Res.*, 24, 601–608, 1977.
- Dzyuban, A. N., Krylova, I. N., and Kuznetsova, I. A.: Properties of bacteria distribution and gas regime within the water column of the Baltic Sea in winter, *Oceanology*, 39, 348–351, 1999.
- Elkins, J. W.: Aquatic sources and sinks for nitrous oxide, PhD thesis, Harvard University, Cambridge, MA, 1978.
- Elkins, J. W., Wofsy, S. C., McElroy, M. B., Kolb, C. E., and Kaplan, W. A.: Aquatic sources and sinks for nitrous oxide, *Nature*, 275, 602–606, 1978.
- Emeis, K. C., Brüchert, V., Currie, B., Endler, R., Ferdelman, T., Kiessling, A., Leipe, T., Noli-Peard, K., Struck, U., and Vogt, T.: Shallow gas in shelf sediments of the Namibian coastal upwelling ecosystem, *Cont. Shelf Res.*, 24, 627–642, 2004.
- Farías, L. and Cornejo, M.: Effect of seasonal change in bottom water oxygenation on sediment N-oxide and N<sub>2</sub>O cycling in the coastal upwelling regime off central Chile (36.5° S), *Prog. Oceanogr.*, 75, 561–575, 2007.
- Farías, L., Graco, M., and Ulloa, O.: Temporal variability of nitrogen cycling in continental-shelf sediments of the upwelling ecosystem off Central Chile, *Deep-Sea Res. II*, 51, 2491–2505, 2004.
- Farías, L., Paulmier, A., and Gallegos, M.: Nitrous oxide and N-nutrient cycling in the oxygen minimum zone off northern Chile, *Deep-Sea Res. I*, 54, 164–180, 2007.
- Farías, L., Castro-González, M., Cornejo, M., Charpentier, J., Faúndez, J., Boontanon, N., and Yoshida, N.: Denitrification and nitrous oxide cycling within the upper oxycline of the oxygen minimum zone off the eastern tropical South Pacific, *Limnol. Oceanogr.*, 54, 132–144, 2009a.
- Farías, L., Fernández, C., Faúndez, J., Cornejo, M., and Alcamán, M. E.: Chemolithoautotrophic production mediating the cycling of the greenhouse gases N<sub>2</sub>O and CH<sub>4</sub> in an upwelling ecosystem, *Biogeosciences*, 6, 3053–3069, doi:10.5194/bg-6-3053-2009, 2009b.
- Fenchel, T., Bernard, C., Esteban, G., Finlay, B. J., Hansen, P. J., and Iversen, N.: Microbial diversity and activity in a Danish fjord with anoxic deep water, *Ophelia*, 43, 45–100, 1995.
- Fox, L. E., Lipschultz, F., Kerkhof, L., and Wofsy, S. C.: A chemical survey of the Mississippi Estuary, *Estuaries*, 10, 1–12, 1987.
- Galloway, J. N., Dentener, F. J., Capone, D. G., Boyer, E. W., Howarth, R. W., Seitzinger, S. P., Asner, G. P., Cleveland, C. C., Green, P. A., Holland, E. A., Karl, D. M., Michaels, A. F., Porter, J. H., Townsend, A. R., and Vorosmarty, C. J.: Nitrogen cycles: past, present and future, *Biogeochemistry*, 70, 153–226, 2004.
- Garcia, H. E., Locarnini, R. A., Boyer, T. P., and Antonov, J. I.: World Ocean Atlas 2005, Vol. 3: Dissolved Oxygen, Apparent Oxygen Utilization, and Oxygen Saturation. S. Levitus, Ed. NOAA Atlas NESDIS 63, US Government Printing Office, Washington, D.C., 342 pp., 2006.
- Goreau, T. J., Kaplan, W. A., Wofsy, S. C., McElroy, M. B., Valois, F. W., and Watson, S. W.: Production of NO<sub>2</sub> and N<sub>2</sub>O by nitrifying bacteria at reduced concentrations of oxygen, *Appl. Environ. Microbiol.*, 40, 526–532, 1980.
- Grantham, B. A., Chan, F., Nielsen, K. J., Fox, D. S., Barth, J. A., Huyer, A., Lubchenco, J., and Menge, B. A.: Upwelling-driven

- nearshore hypoxia signals ecosystem and oceanographic changes in the northeast Pacific, *Nature*, 429, 749–754, 2004.
- Gruber, N.: The dynamics of the marine nitrogen cycle and its influence on atmospheric CO<sub>2</sub>, in: *The Ocean Carbon Cycle and Climate*, edited by: Follows, M. and Oguz, T., NATO ASI Series, Kluwer Academic, Dordrecht, 97–148, 2004.
- Hagy, J. D., Boynton, W. R., Keefe, C. W., and Wood, K. V.: Hypoxia in Chesapeake Bay, 1950–2001: Long-term change in relation to nutrient loading and river flow, *Estuaries*, 27, 634–658, 2004.
- Hashimoto, S., Gojo, K., Hikota, S., Sendai, N., and Otsuki, A.: Nitrous oxide emissions from coastal waters in Tokyo Bay, *Mar. Environ. Res.*, 47, 213–223, 1999.
- Hashimoto, L. K., Kaplan, W. A., Wofsy, S. C., and McElroy, M. B.: Transformation of fixed nitrogen and N<sub>2</sub>O in the Cariaco Trench, *Deep-Sea Res. A*, 30, 575–590, 1983.
- Helly, J. J. and Levin, L. A.: Global distribution of naturally occurring marine hypoxia on continental margins, *Deep-Sea Res. I*, 51, 1159–1168, 2004.
- IPCC: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, UK and New York, NY, USA, 2007.
- Jayakumar, D. A.: Biogeochemical cycling of methane and nitrous oxide in the northern Indian Ocean. PhD Thesis, Goa University, Goa, India, 1999.
- Jayakumar, D. A., Naqvi, S. W. A., Narvekar, P. V., and George, M. D.: Methane in coastal and offshore waters of the Arabian Sea, *Mar. Chem.*, 74, 1–13, 2001.
- Jickells, T.: The role of air-sea exchange in the marine nitrogen cycle, *Biogeosciences*, 3, 271–280, doi:10.5194/bg-3-271-2006, 2006.
- Joos, F., Plattner, G. -K., Stocker, T. F., Körtzinger, A., and Wallace, D. W. R.: Trends in marine dissolved oxygen: Implications for ocean circulation changes and the carbon budget, *EOS-Trans. Amer. Geophys. Union*, 84, doi:10.1029/2003EO210001, 2003.
- Karisiddaiah, S. M. and Veerayya, M.: Methane-bearing shallow gas-charged sediments in the eastern Arabian Sea: a probable source for greenhouse gas, *Cont. Shelf Res.*, 14, 1361–1370, 1994.
- Karl, D. M. and Tilbrook, B. D.: Production and transport of methane in oceanic particulate organic matter, *Nature*, 368, 732–734, 1994.
- Karl, D. M., Beversdorf, L., Bjoerkman, K. M., Church, M. J., Martinez, A., DeLong, E. F. Aerobic production of methane in the sea, *Nature Geosci.*, 1, 473–478, 2008.
- Kelley, C.: Methane oxidation potential in the water column of two diverse coastal marine sites, *Biogeochemistry*, 65, 105–120, 2003.
- Kelley, C. A. and Jeffrey, W. H.: Dissolved methane concentration profiles and air-sea fluxes from 41° S to 27° N, *Global Biogeochem. Cycles*, 16, 1040, doi:10.1029/2001GB001809, 2002.
- Kemp, W. M., Testa, J. M., Conley, D. J., Gilbert, D., and Hagy, J. D.: Temporal responses of coastal hypoxia to nutrient loading and physical controls, *Biogeosciences*, 6, 2985–3008, doi:10.5194/bg-6-2985-2009, 2009.
- Kessler, J. D., Reeburgh, W. S., Southon, J., and Varela, R.: Fossil methane source dominates Cariaco Basin water column methane geochemistry, *Geophys. Res. Lett.*, 32, L12609, doi:10.1029/2005GL022984, 2005.
- Kessler, J. D., Reeburgh, W. S., Southon, J., Seifert, R., Michaelis, W., and Tyler, S. C.: Basin-wide estimates of the input of methane from seeps and clathrates to the Black Sea, *Earth Planet. Sci. Lett.*, 243, 366–375, 2006a.
- Kessler, J. D., Reeburgh, W. S., and Tyler, S. C.: Controls on methane concentration and stable isotope ( $\delta^2\text{H-CH}_4$  and  $\delta^{13}\text{C-CH}_4$ ) distributions in the water columns of the Black Sea and Cariaco Basin, *Global Biogeochem. Cycles*, 20, GB4004, doi:10.1029/2005GB002571, 2006b.
- Kessler, J. D., Reeburgh, W. S., Valentine, D. L., Kinnaman, F. S., Peltzer, E. T., Brewer, P. G., Southon, J., and Tyler, S. C.: A survey of methane isotope abundance ( $^{14}\text{C}$ ,  $^{13}\text{C}$ ,  $^2\text{H}$ ) from five nearshore marine basins that reveals unusual radiocarbon levels in subsurface waters, *J. Geophys. Res.*, 113, C12021, doi:10.1029/2008JC004822, 2008.
- Kim, K.-R. and Craig, H.: Two isotope characterisation of N<sub>2</sub>O in the Pacific Ocean and constraints on its origin in deep water, *Nature*, 347, 58–61, 1990.
- Kock, A., Gebhardt, S., and Bange, H. W.: Methane emissions from the upwelling area off Mauritania (NW Africa), *Biogeosciences*, 5, 1119–1125, doi:10.5194/bg-5-1119-2008, 2008.
- Konavalov, S. K., Murray, J. W., and Luther III, G. W.: Basic processes of Black Sea biogeochemistry, *Oceanography*, 18, 24–35, 2005.
- Kuypers, M. M. M., Lavik, G., Wöbken, D., Schmid, M., Fuchs, B. M., Amann, R., Jørgensen, B. B., and Jetten, M. S. M.: Massive nitrogen loss from the Benguela upwelling system through anaerobic ammonium oxidation, *Proc. Nat. Acad. Sci. USA*, 102, 6478–6483, 2005.
- Lam, P., Lavik, G., Jensen, M. M., van de Vossenberg, J., Schmid, M., Wöbken, D., Gutierrez, D., Amann, R., Jetten, M. S. M., and Kuypers, M. M. M.: Revising the nitrogen cycle in the Peruvian oxygen minimum zone, *Proc. Nat. Acad. Sci. USA*, 106, 4752–4757, 2009.
- Lavik, G., Stührmann, T., Brüchert, V., van der Plas, A., Mohrholz, V., Lam, P., Mußmann, M., Fuchs, B. M., Amann, R., Lass, U., Kuypers, M. M. M.: Detoxification of sulphidic African shelf waters by blooming chemolithotrophs, *Nature*, 457, 581–584, 2009.
- Law, C. S. and Owens, N. J. P.: Significant flux of atmospheric nitrous oxide from the northwest Indian Ocean, *Nature*, 346, 826–828, 1990.
- Levin, L. A., Ekau, W., Gooday, A. J., Jorissen, F., Middelburg, J. J., Naqvi, S. W. A., Neira, C., Rabalais, N. N., and Zhang, J.: Effects of natural and human-induced hypoxia on coastal benthos, *Biogeosciences*, 6, 2063–2098, doi:10.5194/bg-6-2063-2009, 2009.
- Li, D. J., Zhang, J., Huang, D. J., Wu, Y., and Liang, J.: Oxygen depletion off the Changjiang (Yangtze River) Estuary, *Sci. China*, 45D, 1137–1146, 2002.
- Lilley, M. D., Baross, J. A., and Gordon, L. I.: Dissolved hydrogen and methane in Saanich Inlet, British Columbia, *Deep-Sea Res.*, 29, 1471–1484, 1982.
- Luyten, J. R., Pedlosky, J., and Stommel, H.: The ventilated thermocline, *J. Phys. Oceanogr.*, 13, 292–309, 1983.
- Martens, C. S. and Klump, J. V.: Biogeochemical cycling in an organic-rich coastal marine basin – I. Methane sediment-water

- exchange processes, *Geochim. Cosmochim. Acta*, 44, 471–490, 1980.
- McGinnis, D. F., Greinert, J., Artemov, Y., Beaubien, S. E., and Wüest, A.: Fate of rising methane bubbles in stratified waters: How much methane reaches the atmosphere?, *J. Geophys. Res.*, 111, C09007, doi:10.1029/2005JC003183, 2006.
- Milliman, J. D. and Meade, R. H.: Worldwide delivery of river sediment to the oceans, *J. Geol.*, 91, 1–21, 1983.
- Molina, V. and Farías, L.: Aerobic ammonium oxidation in the oxycline and oxygen minimum zone of the eastern tropical South Pacific off northern Chile (20° S), *Deep-Sea Res. II*, 56, 1032–1041, 2009.
- Molina, V., Farías, L., Eissler, Y., Cuevas, L. A., Morales, C. E., and Escribano, R.: Ammonium cycling under a strong oxygen gradient associated with the Oxygen Minimum Zone off northern Chile (~23° S), *Mar. Ecol. Prog. Ser.*, 288, 35–43, 2005.
- Monteiro, P. M. S., van der Plas, A., Mohrholz, V., Mabilille, E., Pascall, A., and Joubert, W.: Variability of natural hypoxia and methane in a coastal upwelling system: Oceanic physics or shelf biology?, *Geophys. Res. Lett.*, 33, L16614, doi:10.1029/2006GL026234, 2006.
- Monteiro, P. M. S., van der Plas, A. K., Melice, J.-L., and Florenchie, P.: Interannual hypoxia variability in a coastal upwelling system: Ocean-shelf exchange, climate and ecosystem-state implications, *Deep-Sea Res. I*, 55, 435–450, 2008.
- Morrison, J. M., Codispoti, L. A., Gaurin, S., Jones, B., Manghanani, V., and Zheng, Z.: Seasonal variation of hydrographic and nutrient fields during the US JGOFS Arabian Sea Process Study, *Deep-Sea Res. II*, 45, 2053–2101, 1998.
- Naqvi, S. W. A.: Some aspects of the oxygen deficient conditions and denitrification in the Arabian Sea, *J. Mar. Res.*, 45, 1049–1072, 1987.
- Naqvi, W. A.: Geographical extent of denitrification in the Arabian Sea in relation to some physical processes, *Oceanolog. Acta*, 14, 281–290.
- Naqvi, S. W. A.: Denitrification processes in the Arabian Sea, *Proc. Indian Acad. Sci. (Earth & Planet Sci.)*, 103, 279–300, 1994.
- Naqvi, S. W. A. and Noronha, R. J.: Nitrous oxide in the Arabian Sea, *Deep-Sea Res.*, 38, 871–890, 1991.
- Naqvi, S. W. A., Jayakumar, D. A., Nair, M., Kumar, M. D., and George, M. D.: Nitrous oxide in the western Bay of Bengal, *Mar. Chem.*, 47, 269–278, 1994.
- Naqvi, S. W. A., Yoshinari, T., Jayakumar, D. A., Altabet, M. A., Narvekar, P. V., Devol, A. H., Brandes, J. A., and Codispoti, L. A.: Budgetary and biogeochemical implications of N<sub>2</sub>O isotope signatures in the Arabian Sea, *Nature*, 394, 462–464, 1998.
- Naqvi, S. W. A., Naik, H., and Narvekar, P. V.: The Arabian Sea, in: *Biogeochemistry of Marine Systems*, edited by: Black, K. and Shimmield, G., Sheffield Academic Press, Sheffield, 156–206, 2003.
- Naqvi, S. W. A., Jayakumar, D. A., Narvekar, P. V., Naik, H., Sarma, V. V. S. S., D'Souza, W., Joseph, S., and George, M. D.: Increased marine production of N<sub>2</sub>O due to intensifying anoxia on the Indian continental shelf, *Nature*, 408, 346–349, 2000.
- Naqvi, S. W. A., Bange, H. W., Gibb, S. W., Goyet, C., Hatton, A. D., and Upstill-Goddard, R. C.: Biogeochemical ocean-atmosphere transfers in the Arabian Sea, *Prog. Oceanogr.*, 65, 116–144, 2005.
- Naqvi, S. W. A., Narvekar, P. V., and Desa, E.: Coastal biogeochemical processes in the North Indian Ocean, in: *The Sea*, vol. 14, edited by: Robinson, A., and Brink, K., Harvard University Press, 723–780, 2006a.
- Naqvi, S. W. A., Naik, H., Jayakumar, D. A., Shailaja, M. S., and Narvekar, P. V.: Seasonal oxygen deficiency over the western continental shelf of India, in: *Past and Present Water Column Anoxia*, edited by: Neretin, L. N., NATO Science Series, IV. Earth and Environmental Sciences, vol. 64, Springer, Dordrecht, 195–224, 2006b.
- Naqvi, S. W. A., Naik, H., Pratihary, A., D'Souza, W., Narvekar, P. V., Jayakumar, D. A., Devol, A. H., Yoshinari, T., and Saino, T.: Coastal versus open-ocean denitrification in the Arabian Sea, *Biogeosciences*, 3, 621–633, doi:10.5194/bg-3-621-2006, 2006c.
- Naqvi, S. W. A., Naik, H., Jayakumar, D. A., Pratihary, A., Narvekar, G., Kurian, S., Agnihotri, R., Shailaja, M. S., and Narvekar, P. V.: Seasonal anoxia over the western Indian continental shelf, in: *Indian Ocean: Biogeochemical Processes and Ecological Variability*, edited by: Wiggert, J. D., Hood, R. R., Naqvi, S. W. A., Brink, K. H., and Smith, S. L., *Geophys. Monogr. Ser.*, 185, AGU, Washington, D.C., 333–345, 2009.
- Naqvi, S. W. A., Naik, H., D'Souza, W., Narvekar, P. V., Paropkari, A. L., and Bange, H. W.: Carbon and nitrogen fluxes in the North Indian Ocean, in: *Carbon and nutrient fluxes in continental margins: A global synthesis*, edited by: Liu, K. K., Atkinson, L., Quinones, R., and Talaue-McManus, L., Springer-Verlag, New York, 180–192, 2010a.
- Naqvi, S. W. A., Moffett, J. W., Gauns, M. U., Narvekar, P. V., Pratihary, A. K., Naik, H., Shenoy, D. M., Jayakumar, D. A., Goepfert, T. J., Patra, P. K., Al-Azri, A., and Ahmed, S. I.: The Arabian Sea as a high-nutrient, low-chlorophyll region during the late Southwest Monsoon, *Biogeosciences*, 7, 2091–2100, doi:10.5194/bg-7-2091-2010, 2010b.
- Neretin, L. N. (Ed.): *Past and Present Water Column Anoxia*, NATO Science Series, IV. Earth and Environmental Sciences, vol. 64, Springer, Dordrecht, 541 pp., 2006.
- Nevison, C. D., Weiss, R. F., and Erickson III, D. J.: Global oceanic emissions of nitrous oxide, *J. Geophys. Res.*, 100, 15809–15820, 1995.
- Nevison C. D., Lueker, T. J., and Weiss, R. F.: Quantifying the nitrous oxide source from coastal upwelling, *Global Biogeochem. Cycles*, 18, GB1018, doi:10.1029/2003GB002110, 2004.
- Nicholls, J. C., Davies, C. A., and Trimmer, M.: High-resolution profiles and nitrogen isotope tracing reveal a dominant source of nitrous oxide and multiple pathways of nitrogen gas formation in the central Arabian Sea, *Limnol. Oceanogr.*, 52, 156–168, 2007.
- Oremland, R. S.: Methanogenic activity in plankton samples and fish intestines – mechanism for in situ methanogenesis in oceanic surface waters, *Limnol. Oceanogr.*, 24, 1136–1141, 1979.
- Oschlies, A., Schulz, K. G., Riebesell, U., and Schmittner, A.: Simulated 21st century's increase in oceanic suboxia by CO<sub>2</sub>-enhanced biotic carbon export, *Global Biogeochem. Cycles*, 22, GB4008, doi:10.1029/2007GB003147, 2008.
- Oudot, C., Andrie, C., and Montel, Y.: Nitrous oxide production in the tropical Atlantic Ocean, *Deep-Sea Res.*, 37, 183–202, 1990.
- Owens, N. J. P., Law, C. S., Mantoura, R. F. C., Burkill, P. H., and Lewellyn, C. A.: Methane flux to the atmosphere from the Arabian Sea, *Nature*, 354, 293–296, 1991.
- Paulmier, A. and Ruiz-Pino, D.: Oxygen minimum zones (OMZs)

- in the modern ocean, *Prog. Oceanogr.*, 80, 113–128, 2009.
- Paulmier A., Ruiz-Pino, D., Garçon, V., and Farias, L.: Maintaining of the Eastern South Pacific oxygen minimum zone (OMZ) off Chile, *Geophys. Res. Lett.*, 33, L20601, doi:10.1029/2006GL026801, 2006.
- Pierotti, D. and Rasmussen, R. A.: Nitrous oxide measurements in the eastern tropical Pacific Ocean, *Tellus*, 32, 56–72, 1980.
- Rabalais, N. N. and Turner, R. E.: Oxygen depletion in the Gulf of Mexico adjacent to the Mississippi River, in: *Past and Present Marine Water Column Anoxia*, edited by: Neretin, L. N., NATO Science Series, IV. Earth and Environmental Sciences, vol. 64, Springer, Dordrecht, 225–245, 2006.
- Rabalais, N. N., Turner, R. E., Sen Gupta, B. K., Platon, E., and Parsons, M. L.: Sediments tell the history of eutrophication and hypoxia in the Northern Gulf of Mexico, *Ecol. Appl.*, 17, S129–S143, 2007.
- Rabalais, N. N., Díaz, R. J., Levin, L. A., Turner, R. E., Gilbert, D., and Zhang, J.: Dynamics and distribution of natural and human-caused hypoxia, *Biogeosciences*, 7, 585–619, doi:10.5194/bg-7-585-2010, 2010.
- Reeburgh, W. S.: Methane consumption in Cariaco Trench waters and sediments, *Earth Planet. Sci. Lett.*, 28, 337–344, 1976.
- Reeburgh, W. S.: Oceanic methane biogeochemistry, *Chem. Rev.*, 107, 486–513, 2007.
- Reeburgh, W. S., Ward, B. B., Wahlen, S. C., Sandbeck, K. A., Kilpatrick, K. A., and Kerkhof, L. J.: Black-Sea methane geochemistry, *Deep-Sea Res.*, 38, S1189–S1210, 1991.
- Rehder, G., Keir, R. S., Suess, E., and Rhein, M.: Methane in the northern Atlantic controlled by microbial oxidation and atmospheric history, *Geophys. Res. Lett.*, 26, 587–590, 1999.
- Rehder, G., Collier, R. W., Heeschen, K., Kosro, P. M., Barth, J., and Suess, E.: Enhanced marine CH<sub>4</sub> emissions to the atmosphere off Oregon caused by coastal upwelling, *Global Biogeochem. Cycles*, 16, 1731, doi:10.1029/2000GB001391, 2002.
- Revsbech, N. P., Larsen, L. H., Gundersen, J., Dalsgaard, T., Ulløe, O., and Thamdrup, B.: Determination of ultra-low oxygen concentrations in oxygen minimum zones by the STOX sensor, *Limnol. Oceanogr. Methods*, 7, 371–381, 2009.
- Rhee, T. S., Kettle, A. J., and Andreae, M. O.: Methane and nitrous oxide emissions from the ocean: A reassessment using basin-wide observations in the Atlantic, *J. Geophys. Res.*, 114, D12304, doi:10.1029/2008JD011662, 2009.
- Rönnner, U.: Distribution, production and consumption of nitrous oxide in the Baltic Sea, *Geochim. Cosmochim. Acta*, 47, 2179–2188, 1983.
- Sansone, F. J., Popp, B. N., Gasc, A., Graham, A. W., and Rust, T. M.: Highly elevated methane in the eastern tropical North Pacific and associated isotopically enriched fluxes to the atmosphere, *Geophys. Res. Lett.*, 28, 4567–4570, 2001.
- Sansone, F. J., Graham, A. W., and Berelson, W. M.: Methane along the western Mexican margin, *Limnol. Oceanogr.*, 49, 2242–2255, 2004.
- Schmale, O., Greinert, J., and Rehder, G.: Methane emission from high-intensity marine gas seeps in the Black Sea into the atmosphere, *Geophys. Res. Lett.*, 32, L07609, doi:10.1029/2004GL021138, 2005.
- Schmaljohann, R.: Methane dynamics in the sediment and water column of Kiel Harbour (Baltic Sea), *Mar. Ecol. Prog. Ser.*, 131, 262–273, 1996.
- Schweiger, B.: Messung von NH<sub>2</sub>OH in ausgewählten Seegebieten, Diploma thesis, Kiel University, Kiel, 115 pp., 2006.
- Schweiger, B., Hansen, H. P., and Bange, H. W.: A time series of hydroxylamine (NH<sub>2</sub>OH) in the southwestern Baltic Sea, *Geophys. Res. Lett.*, 34, L24608, doi:10.1029/2007GL031086, 2007.
- Scranton, M. I. and Brewer, P. G.: Occurrence of methane in the near surface waters of the western subtropical North Atlantic, *Deep-Sea Res.*, 24, 127–138, 1977.
- Scranton, M. I. and Farrington, J. W.: Methane production in the waters off Walvis Bay, *J. Geophys. Res.*, 82, 4947–4953, 1977.
- Scranton, M. I. and McShane, K.: Riverine sources of methane to the southern bight of the North Sea, *Cont. Shelf Res.*, 11, 37–52, 1991.
- Seitzinger, S. P. and Kroeze, C.: Global distribution of nitrous oxide production and N inputs in freshwater and coastal marine ecosystems, *Global Biogeochem. Cycles*, 12, 93–113, 1998.
- Seitzinger, S. P., Kroeze, C., Bouwman, A. E., Caraco, N., Dentener, F., Styles, R. V.: Global patterns of dissolved inorganic and particulate nitrogen inputs to coastal systems: Recent conditions and future predictions, *Estuaries*, 25, 640–655, 2002.
- Senga, Y., Mochida, K., Fukumori, R., Okamoto, N., and Seike, Y.: N<sub>2</sub>O accumulation in estuarine and coastal sediments: The influence of H<sub>2</sub>S on dissimilatory nitrate reduction, *Estuar. Coast. Shelf. Sci.*, 67, 231–238, 2006.
- Shaffer, G., Olsen, S. M., and Pedersen, J. O. P.: Long-term ocean oxygen depletion in response to carbon dioxide emissions from fossil fuels, *Nature Geosci.*, 2, 105–109, 2009.
- Shaffer, G., Hormazabal, S., Pizarro, O., and Salinas, S.: Seasonal and interannual variability of currents and temperature off central Chile, *J. Geophys. Res.*, 104, 29951–29961, 1999.
- Shailaja, M. S., Narvekar, P. V., Alagarsamy, R., and Naqvi, S. W. A.: Nitrogen transformations as inferred from the activities of key enzymes in the Arabian Sea oxygen minimum, *Deep-Sea Res. I*, 53, 960–970, 2006.
- Sigman, D. M., Robinson, R., Knapp, A. N., van Geen, A., McCorkle, D. C., Brandes, J. A., and Thunell, R. C.: Distinguishing between water column and sedimentary denitrification in the Santa Barbara Basin using the stable isotopes of nitrate, *Geochem. Geophys. Geosyst.*, 4, 1040, doi:10.1029/2002GC000384, 2003.
- Smith, S. V., Swaney, D. P., Talaue-McManus, L., Bartley, J. D., Sandhei, P. T., McLaughlin, C. J., Dupra, V. C., Crossland, C. J., Buddemeier, R. W., Maxwell, B. A., and Wulff, F.: Humans, hydrology, and the distribution of inorganic nutrient loading to the ocean, *BioScience*, 53, 235–245, 2003.
- Sobarzo, M. and Djurfeldt, L.: Coastal upwelling process on a continental shelf limited by submarine canyons, Concepción, central Chile, *J. Geophys. Res.*, 109, C12012, doi:10.1029/2004JC002350, 2004.
- Solomon, E. A., Kastner, M., MacDonald, I. R., and Leifer, I.: Considerable methane fluxes to the atmosphere from hydrocarbon seeps in the Gulf of Mexico, *Nature Geosci.*, 2, 561–565, 2009.
- Stramma, L., Johnson, G. C., Sprintall, J., and Mohrholz, V.: Expanding oxygen-minimum zones in the tropical oceans, *Science*, 320, 655–658, 2008.
- Strub, P. T., Mesias, J. M., Montecinos, V., Rutllant, J., and Salinas, S.: Coastal ocean circulation off western south America, in: *The Sea*, vol. 11, edited by: Robinson, A. R. and Brink, K. H., John Wiley & Sons, 273–313, 1998.

- Su, J. L.: Circulation dynamics of the China Seas north of 18° N, in: *The Sea*, vol. 11, edited by: Robinson, A. R. and Brink, K. H., John Wiley & Sons, New York, 483–505, 1998.
- Swarzenski, P. W., Campbell, P. L., Osterman, L. E., and Poore, R. Z.: A 1000-year sediment record of recurring hypoxia off the Mississippi River: The potential role of terrestrially-derived organic matter inputs, *Mar. Chem.*, 109, 130–142, 2008.
- Swinerton, J. W. and Lamontagne, R. A.: Oceanic distribution of low-molecular-weight hydrocarbons – Baseline measurements, *Environ. Sci. Technol.*, 8, 657–663, 1974.
- Tilbrook, B. D. and Karl, D. M.: Methane sources, distributions and sinks from California coastal waters to the oligotrophic North Pacific gyre, *Mar. Chem.*, 49, 51–64, 1995.
- Treude, T., Krüger, M., Boetius, A., and Jørgensen, B. B.: Environmental control on anaerobic oxidation of methane in gassy sediments of Eckernförde Bay (German Baltic), *Limnol. Oceanogr.*, 50, 1771–1786, 2005.
- Tsunogai, S., Watanabe, S., and Sato, T.: Is there a “continental shelf pump” for the absorption of atmospheric CO<sub>2</sub>, *Tellus, Ser. B*, 51, 701–712, 1999.
- Upstill-Goddard, R. C., Barnes, J., and Owens, N. J. P.: Nitrous oxide and methane during the 1994 SW monsoon in the Arabian Sea/northwestern Indian Ocean, *J. Geophys. Res.*, 104, 30067–30084, 1999.
- van der Plas, A. K., Monteiro, P. M. S., and Pascall, A.: Cross-shelf biogeochemical characteristics of sediments in the central Benguela and their relationship to overlying water column hypoxia, *Afr. J. Mar. Sci.*, 29, 37–47, 2007.
- van der Star, W. R. L.: Growth and Metabolism of Anammox Bacteria, *Sieca Repro*, Delft, Netherlands, 153 pp., 2008.
- Walker, J. T., Stow, C. A., and Geron, C.: Nitrous oxide emissions from the Gulf of Mexico Hypoxic Zone, *Environ. Sci. Technol.*, 44, 1617–1623, 2010.
- Walter, S., Bange, H. W., and Wallace, D. W. R.: Nitrous oxide in the surface layer of the tropical North Atlantic Ocean along a west to east transect, *Geophys. Res. Lett.*, 31, L23S07, doi:10.1029/2004GL019937, 2004.
- Walter, S., Breitenbach, U., Bange, H. W., Nausch, G., and Wallace, D. W. R.: Distribution of N<sub>2</sub>O in the Baltic Sea during transition from anoxic to oxic conditions, *Biogeosciences*, 3, 557–570, doi:10.5194/bg-3-557-2006, 2006.
- Ward, B. B.: The subsurface methane maximum in the Southern California Bight, *Cont. Shelf Res.*, 12, 735–752, 1992.
- Ward, B. B., Kilpatrick, K. A., Wopat, A. E., Minnich, E. C., and Lidstrom, M. E.: Methane oxidation in Saanich Inlet during summer stratification, *Cont. Shelf Res.*, 9, 65–75, 1989.
- Ward, B. B., Kilpatrick, K. A., Novelli, P. C., and Scranton, M. I.: Methane oxidation and methane fluxes in the ocean surface layer and deep anoxic waters, *Nature*, 327, 226–229, 1987.
- Ward, B. B., Tuit, C. B., Jayakumar, A., Rich, J. J., Moffett, J. W., and Naqvi, S. W. A.: Organic carbon, not copper, controls denitrification in oxygen minimum zones of the ocean, *Deep-Sea Res. I*, 55, 1672–1683, 2008.
- Weeks, S. J., Currie, B., and Bakun, A.: Satellite imaging: Massive emissions of toxic gas in the Atlantic, *Nature*, 415, 493–494, 2002.
- Welsh, D. T., Castadelli, G., Bartoli, M., Poli, D., Careri, M., de Wit, R., and Viaroli, P.: Denitrification in an intertidal seagrass meadow, a comparison of <sup>15</sup>N-isotope and acetylene-block techniques: dissimilatory nitrate reduction to ammonia as a source of N<sub>2</sub>O? *Mar. Biol.*, 139, 1029–1036, 2001.
- Westley, M. B., Yamagishi, H., Popp, B. B., and Yoshida, N.: Nitrous oxide cycling in the Black Sea inferred from stable isotope and isotopomer distributions, *Deep-Sea Res. II*, 53, 1802–1816, 2006.
- Wyrtki, K.: *Oceanographic Atlas of the International Indian Ocean Expedition*, National Science Foundation, Washington, D.C., 531 pp., 1971.
- Yamagishi H., Westley, M. B., Popp, B. N., Toyoda, S., Yoshida, N., Watanabe, S., Koba, K., and Yamanaka, Y.: Role of nitrification and denitrification on the nitrous oxide cycle in the eastern tropical North Pacific and Gulf of California, *J. Geophys. Res.*, 112, G02015, doi:10.1029/2006JG000227, 2007.
- Yang, S. L., Zhao, Q. Y., and Belkin, I. M.: Temporal variation in the sediment load of the Yangtze River and the influences of the human activities, *J. Hydrol.*, 263, 56–71, 2002.
- Yin, K. D., Lin, Z. F., and Ke, Z. Y.: Temporal and spatial distribution of dissolved oxygen in the Pearl River Estuary and adjacent coastal waters, *Cont. Shelf Res.*, 24, 1935–1948, 2004.
- Yoshinari, T.: Nitrous oxide in the sea, *Mar. Chem.*, 4, 189–202, 1976.
- Yoshinari, T., Altabet, M. A., Naqvi, S. W. A., Codispoti, L. A., Jayakumar, A., Kuhland, M., and Devol, A. H.: Nitrogen and oxygen isotopic composition of N<sub>2</sub>O from suboxic waters of the eastern tropical North Pacific and the Arabian Sea – measurement by continuous-flow isotope ratio monitoring, *Mar. Chem.*, 56, 253–264, 1997.
- Zhang, G. L., Zhang, J., Kang, Y. B., and Liu, S. M.: Distribution and fluxes of methane in the East China Sea and Yellow Sea in spring, *J. Geophys. Res.*, 109, C07011, doi:10.1029/2004JC002268, 2004.
- Zhang, G. L., Zhang, J., Liu, S. M., Ren, J. L., Xu, J., and Zhang, F.: Methane in the Changjiang (Yangtze River) Estuary and its adjacent marine area: Riverine input, sediment release and atmospheric fluxes, *Biogeochemistry*, 91, 71–84, 2008a.
- Zhang, G. L., Zhang, J., Ren, J. L., Li, J. B., and Liu, S. M.: Distribution and sea-to-air fluxes of methane and nitrous oxide in the North East China Sea in summer, *Mar. Chem.*, 110, 42–55, 2008b.